

Product datasheet

Anti-Aggrecan antibody [6-B-4] ab3778

★★★★☆ 5 Abreviews 32 References 3 图像

概述

产品名称	Anti-Aggrecan抗体[6-B-4]
描述	小鼠单克隆抗体[6-B-4] to Aggrecan
宿主	Mouse
特异性	Recognises the linear amino acid sequence 394EPEEPFTFAPEI406 present in the interglobular domain (IGD) of human and bovine aggrecan (in both intact and matrix protease degraded aggrecan) but not pig and several other animals.
经测试应用	适用于: ELISA, IHC-P, IHC-Fr, WB, ICC/IF
种属反应性	与反应: Mouse, Cow, Human
免疫原	Synthetic peptide: EPEEPFTFAPEI , corresponding to amino acids 413-424 of Human Aggrecan.

[Run BLAST with](#) [Run BLAST with](#)

常规说明 Samples are usually deglycosylated using 0.01 Units Chondroitinase ABC (Sigma), 0.01 Units Keratanase (Seikagaku) and 0.0001 Units Keratanase II (Seikagaku) per 10µg S-GAG of non-deglycosylated aggrecan for optimal epitope recognition in SDS-PAGE and immunohistochemistry – see Little et al. and Caterson et al. Diluted Technomouse culture supernatant.

This antibody detects aggrecan metabolites (intact or matrix protease-catabolised) in human synovial fluid samples.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	0.2M Tris/HCl pH 7.4, 1-10% foetal calf serum, 0.02% sodium azide
纯度	Tissue culture supernatant
Primary antibody说明	This antibody detects aggrecan metabolites (intact or matrix protease-catabolised) in human synovial fluid samples.
克隆	单克隆

克隆编号	6-B-4
骨髓瘤	x63-Ag8.653
同种型	IgG1
轻链类型	kappa

应用

Our [Abpromise guarantee](#) covers the use of **ab3778** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
IHC-P	★★★★☆	Use at an assay dependent concentration.
IHC-Fr	★★★★☆	Use at an assay dependent concentration.
WB		1/100. The antibody will see a large range of aggrecan metabolites/catabolites ranging from greater than 250kD down to 50-60kDa.
ICC/IF	★★★★☆	Use at an assay dependent concentration. PubMed: 21192818

靶标

相关性 Aggrecan is a member of a family of large, aggregating proteoglycans (also including versican, brevican and neurocan) which is found in articular cartilage. Aggrecan is composed of three major domains: G1, G2, and G3. Between the G1 and G2 domains there is an interglobulin region (IGD). The IGD region is the major site of cleavage by specific proteases like metalloproteinases (MMPs) and aggrecanase. Aggrecan cleavage has been associated with a number of degenerative diseases including rheumatoid arthritis and osteoarthritis. There is evidence that this family of proteoglycans modulates cell adhesion, migration, and axonal outgrowth in the CNS.

细胞定位 Secreted; extracellular matrix.


图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aggrecan antibody [6-B-4] (ab3778)

This image is courtesy of an Abreview submitted by Ms Elizabeth Chlipala

ab3778 at 1/100 staining human articular cartilage by IHC-P. The tissue was formaldehyde fixed and blocked with a serum free protein block. An enzymatic antigen retrieval step was performed and the tissue was then incubated with the antibody for 1 hour. An HRP conjugated goat polyclonal antibody was used as the secondary.




Immunocytochemistry/ Immunofluorescence - Anti-Aggrecan antibody [6-B-4] (ab3778)

Image from Berg Vet et al., Arthritis Res Ther. 2010;12(6):R228. Epub 2010 Dec 30. Fig 1.; doi:10.1186/ar3215; 30 December 2010, Arthritis Research & Therapy 2010, 12:R228

Immunofluorescence analysis of Human articular chondrocytes, staining Aggrecan with ab3778.

Cells were fixed with paraformaldehyde and blocked with 1% BSA for 1 hour. Cells were incubated with primary antibody overnight at 4°C. An AlexaFluor®488-conjugated rabbit anti-mouse IgG was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence -

Aggrecan antibody [6-B-4] (ab3778)

Image from Limin Wang, MS et al, TISSUE ENGINEERING: Part A Volume 15, Number 5, 2009, Fig 5.

ab3778 used at a 1/50 dilution staining Aggrecan in human umbilical cord matrix mesenchymal stromal cells by Immunocytochemistry/ Immunofluorescence. Frozen sections of the 3D constructs (10µm) (n=2) were rehydrated with PBS for 5 minutes, and endogenous peroxidase activity was inhibited using 1% hydrogen peroxide in methanol for 30 minutes. The sections were then blocked in 3% horse serum for 20 minutes and incubated with a primary antibody, ab3778, for 1 hour at a 1/50 dilution. After primary antibody incubation the sections were incubated with a streptavidin linked horse anti-mouse IgG secondary antibody for 30 minutes. After secondary antibody incubation, the sections were incubated with avidin-biotinylated enzyme complex for 30 minutes, and then VIP substrate (purple color) was applied on sections for 4 minutes.

The image shown is at 4 weeks, 25M.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors