

Product datasheet

Anti-Alpha 1 microglobulin antibody ab47980

2 图像

概述

产品名称	Anti-Alpha 1 microglobulin抗体
描述	兔多克隆抗体to Alpha 1 microglobulin
宿主	Rabbit
经测试应用	适用于: WB, ICC/IF, IHC-P, ELISA, RIA
种属反应性	与反应: Human
免疫原	Human Alpha 1 Microglobulin

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	Preservative: 0.01% Thimerosal (merthiolate) Constituents: 50% Glycerol, PBS, pH 7.5
纯度	Protein G purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab47980** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
WB		Use at an assay dependent concentration. Predicted molecular weight: 39 kDa.
ICC/IF		Use a concentration of 1 - 5 µg/ml.
IHC-P		Use a concentration of 4 µg/ml.
ELISA		Use at an assay dependent concentration.

应用	Ab评论	说明
RIA		Use at an assay dependent concentration.

靶标


功能	<p>Inter-alpha-trypsin inhibitor inhibits trypsin, plasmin, and lysosomal granulocytic elastase. Inhibits calcium oxalate crystallization.</p> <p>Trypstatin is a trypsin inhibitor.</p>
组织特异性	<p>Expressed by the liver and secreted in plasma. Alpha-1-microglobulin occurs in many physiological fluids including plasma, urine, and cerebrospinal fluid. Inter-alpha-trypsin inhibitor is present in plasma and urine.</p>
序列相似性	<p>In the N-terminal section; belongs to the calycin superfamily. Lipocalin family.</p> <p>Contains 2 BPTI/Kunitz inhibitor domains.</p>
翻译后修饰	<p>The precursor is proteolytically processed into separately functioning proteins.</p> <p>3-hydroxykynurenine, an oxidized tryptophan metabolite that is common in biological fluids, reacts with Cys-53, Lys-111, Lys-137, and Lys-149 to form heterogeneous polycyclic chromophores including hydroxanthommatin. The reaction by alpha-1-microglobulin is autocatalytic; the human protein forms chromophore even when expressed in insect and bacterial cells. The chromophore can react with accessible cysteines forming non-reducible thioether cross-links with other molecules of alpha-1-microglobulin or with other proteins such as Ig alpha-1 chain C region 'Cys-352'.</p> <p>Heavy chains are interlinked with bikunin via a chondroitin 4-sulfate bridge to the their C-terminal aspartate.</p> <p>Addition of glycosaminoglycan chondroitin sulfate, allows cross-linking between the different components.</p>
细胞定位	<p>Secreted.</p>

图片



Immunocytochemistry/ Immunofluorescence - Anti-Alpha 1 microglobulin antibody (ab47980)

ICC/IF image of ab47980 stained MCF7 cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab47980, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Alpha 1 microglobulin antibody (ab47980)

ab47980 Microglobulin alpha 1 in human liver. Left panel: with primary antibody at 4 ug/ml. Right panel: isotype control. Sections were stained using an automated system (DAKO Autostainer Plus), at room temperature: sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffers EDTA pH 9.0. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako envision flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors