

Product datasheet

Anti-Androgen Receptor antibody - ChIP Grade ab3509

★★★★☆ 3 Abreviews 3 References 1 图像

概述

产品名称	Anti-Androgen Receptor抗体- ChIP Grade
描述	兔多克隆抗体to Androgen Receptor - ChIP Grade
宿主	Rabbit
特异性	This antibody detects Androgen Receptor. It does not detect estrogen, progesterone, or glucocorticoid receptors. Immunohistochemical staining of AR in rat brain results in intense nuclear staining.
经测试应用	适用于: ICC, IHC-Fr, IP, WB, ICC/IF, ChIP, IHC-P
种属反应性	与反应: Mouse, Rat, Sheep, Hamster, Human, Pig, Non human primates
免疫原	Fusion protein corresponding to Human Androgen Receptor aa 331-572.
阳性对照	rat brain

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituent: 99% PBS
纯度	Whole antiserum
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab3509** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ICC		Use at an assay dependent concentration.

应用	Ab评论	说明
IHC-Fr		1/10 - 1/100.
IP		Use at an assay dependent concentration.
WB	★★★★☆	1/500. Detects a band of approximately 110 kDa (predicted molecular weight: 110 kDa).
ICC/IF		Use at an assay dependent concentration. PubMed: 22850685
ChIP		Use at an assay dependent concentration.
IHC-P		1/10 - 1/100.

靶标

功能	<p>Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Transcription factor activity is modulated by bound coactivator and corepressor proteins. Transcription activation is down-regulated by NR0B2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3.</p> <p>Isoform 3 and isoform 4 lack the C-terminal ligand-binding domain and may therefore constitutively activate the transcription of a specific set of genes independently of steroid hormones.</p>
组织特异性	<p>Isoform 2 is mainly expressed in heart and skeletal muscle (PubMed:15634333). Isoform 3 is expressed by basal and stromal cells of prostate (at protein level) (PubMed:19244107).</p>
疾病相关	<p>Androgen insensitivity syndrome</p> <p>Spinal and bulbar muscular atrophy X-linked 1</p> <p>Defects in AR may play a role in metastatic prostate cancer. The mutated receptor stimulates prostate growth and metastases development despite of androgen ablation. This treatment can reduce primary and metastatic lesions probably by inducing apoptosis of tumor cells when they express the wild-type receptor.</p> <p>Androgen insensitivity, partial</p>
序列相似性	<p>Belongs to the nuclear hormone receptor family. NR3 subfamily.</p> <p>Contains 1 nuclear receptor DNA-binding domain.</p>
结构域	<p>Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain. In the presence of bound steroid the ligand-binding domain interacts with the N-terminal modulating domain, and thereby activates AR transcription factor activity. Agonist binding is required for dimerization and binding to target DNA. The transcription factor activity of the complex formed by ligand-activated AR and DNA is modulated by interactions with coactivator and corepressor proteins. Interaction with RANBP9 is mediated by both the N-terminal domain and the DNA-binding domain. Interaction with EFCAB6/DJBP is mediated by the DNA-binding domain.</p>
翻译后修饰	<p>Sumoylated on Lys-388 (major) and Lys-521. Ubiquitinated. Deubiquitinated by USP26. 'Lys-6' and 'Lys-27'-linked polyubiquitination by RNF6 modulates AR transcriptional activity and specificity.</p> <p>Phosphorylated in prostate cancer cells in response to several growth factors including EGF. Phosphorylation is induced by c-Src kinase (CSK). Tyr-535 is one of the major phosphorylation sites and an increase in phosphorylation and Src kinase activity is associated with prostate</p>

cancer progression. Phosphorylation by TNK2 enhances the DNA-binding and transcriptional activity and may be responsible for androgen-independent progression of prostate cancer. Phosphorylation at Ser-83 by CDK9 regulates AR promoter selectivity and cell growth. Phosphorylation by PAK6 leads to AR-mediated transcription inhibition. Palmitoylated by ZDHHC7 and ZDHHC21. Palmitoylation is required for plasma membrane targeting and for rapid intracellular signaling via ERK and AKT kinases and cAMP generation.


细胞定位

Nucleus. Cytoplasm. Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1.

形式

There are 2 isoforms produced by alternative splicing. Isoform 1 is also known as: AR-B; isoform 2 is known as AR-A or variant AR45.

图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody - ChIP Grade (ab3509)

Immunohistochemistry (Formaline/PFA-fixed paraffin-embedded sections) analysis of human prostate tissue labeling Androgen Receptor with ab3509 at 1/20 (right). Antigen retrieval was performed using 10mM sodium citrate (pH 6.0), microwaved for 8-15 minutes. Tissues were blocked in 3% H₂O₂-methanol for 15 minutes at room temperature. A HRP-conjugated secondary antibody was used. Left - negative control.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors