

## Product datasheet

# Anti-Androgen Receptor antibody - ChIP Grade ab74272

★★★★★ 13 Abreviews 28 References 5 图像

### 概述

产品名称	Anti-Androgen Receptor抗体- ChIP Grade
描述	兔多克隆抗体to Androgen Receptor - ChIP Grade
经测试应用	适用于: ChIP, IP, WB, ICC/IF, IHC-P, IHC-Fr
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide corresponding to Human Androgen Receptor (N terminal). Sequence is propeitary.
阳性对照	IHC-P: Human prostate carcinoma tissue. WB: LnCap cell lysate.

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.60 Preservative: 0.1% Sodium azide Constituents: 1% BSA, PBS
纯度	Protein A purified
克隆	多克隆
同种型	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab74272** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ChIP	★★★★☆	Use at an assay dependent concentration.
IP		Use at an assay dependent concentration. PubMed: 24466341

应用	Ab评论	说明
WB	★★★★★	Use a concentration of 1 µg/ml. Predicted molecular weight: 99 kDa.
ICC/IF	★★★★★	Use at an assay dependent concentration.
IHC-P	★★★★★	1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
IHC-Fr	★★★★★	Use at an assay dependent concentration. PubMed: 21430025

## 靶标

<b>功能</b>	<p>Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Transcription factor activity is modulated by bound coactivator and corepressor proteins. Transcription activation is down-regulated by NR0B2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3.</p> <p>Isoform 3 and isoform 4 lack the C-terminal ligand-binding domain and may therefore constitutively activate the transcription of a specific set of genes independently of steroid hormones.</p>
<b>组织特异性</b>	Isoform 2 is mainly expressed in heart and skeletal muscle (PubMed:15634333). Isoform 3 is expressed by basal and stromal cells of prostate (at protein level) (PubMed:19244107).
<b>疾病相关</b>	<p>Androgen insensitivity syndrome</p> <p>Spinal and bulbar muscular atrophy X-linked 1</p> <p>Defects in AR may play a role in metastatic prostate cancer. The mutated receptor stimulates prostate growth and metastases development despite of androgen ablation. This treatment can reduce primary and metastatic lesions probably by inducing apoptosis of tumor cells when they express the wild-type receptor.</p> <p>Androgen insensitivity, partial</p>
<b>序列相似性</b>	<p>Belongs to the nuclear hormone receptor family. NR3 subfamily.</p> <p>Contains 1 nuclear receptor DNA-binding domain.</p>
<b>结构域</b>	Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain. In the presence of bound steroid the ligand-binding domain interacts with the N-terminal modulating domain, and thereby activates AR transcription factor activity. Agonist binding is required for dimerization and binding to target DNA. The transcription factor activity of the complex formed by ligand-activated AR and DNA is modulated by interactions with coactivator and corepressor proteins. Interaction with RANBP9 is mediated by both the N-terminal domain and the DNA-binding domain. Interaction with EFCAB6/DJBP is mediated by the DNA-binding domain.
<b>翻译后修饰</b>	<p>Sumoylated on Lys-388 (major) and Lys-521. Ubiquitinated. Deubiquitinated by USP26. 'Lys-6' and 'Lys-27'-linked polyubiquitination by RNF6 modulates AR transcriptional activity and specificity.</p> <p>Phosphorylated in prostate cancer cells in response to several growth factors including EGF. Phosphorylation is induced by c-Src kinase (CSK). Tyr-535 is one of the major phosphorylation sites and an increase in phosphorylation and Src kinase activity is associated with prostate cancer progression. Phosphorylation by TNK2 enhances the DNA-binding and transcriptional activity and may be responsible for androgen-independent progression of prostate cancer.</p>

Phosphorylation at Ser-83 by CDK9 regulates AR promoter selectivity and cell growth.  
 Phosphorylation by PAK6 leads to AR-mediated transcription inhibition.  
 Palmitoylated by ZDHHC7 and ZDHHC21. Palmitoylation is required for plasma membrane targeting and for rapid intracellular signaling via ERK and AKT kinases and cAMP generation.


**细胞定位**

Nucleus. Cytoplasm. Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1.

**形式**

There are 2 isoforms produced by alternative splicing. Isoform 1 is also known as: AR-B; isoform 2 is known as AR-A or variant AR45.

**图片**



**Lane 1 :** Anti-Androgen Receptor antibody - ChIP Grade (ab74272) at 1/50 dilution

**Lane 2 :** Anti-Androgen Receptor antibody - ChIP Grade (ab74272) at 1/100 dilution


**Lane 1 :** LNCaP Whole Cell

**Lane 2 :** LNCaP Whole Cell

**Predicted band size :** 99 kDa

**Observed band size :** 110 kDa


Western blot - Anti-Androgen Receptor antibody - ChIP Grade (ab74272)



[ab74272](#) used in ChIP application. Whole cell lysate from human LNCaP cells was used. Cross linking step (X-ChIP) for 10 minutes in 1% formaldehyde. Taqman detection step. Positive control was PSA enhancer. Negative control region, no antibody control, and in the absence of ligand. 25 µl undiluted [ab74272](#) per IP was used.


ChIP - Anti-Androgen Receptor antibody - ChIP Grade (ab74272)

Image courtesy of an anonymous Abreview.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody - ChIP Grade (ab74272)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human prostate carcinoma labelling Androgen Receptor with ab74272 at a dilution of 1/200 for 30mins at room temperature.



Immunocytochemistry/ Immunofluorescence - Androgen Receptor antibody (ab74272)

This image is courtesy of an Abreview by S. Johann.

ab74272 staining Androgen Receptor in mouse neuroblastoma cells by Immunocytochemistry/Immunofluorescence. Cells were fixed in formaldehyde and permeabilized in 0.2% Triton X-100 prior to blocking in 3% BSA for 1 hour. The primary antibody was diluted 1/200 and incubated with the sample for 12 hours at 4°C. The secondary antibody was Alexa Fluor® 594-conjugated goat anti-rabbit polyclonal, diluted 1/400.



Immunohistochemistry (Frozen sections) - Anti-Androgen Receptor antibody - ChIP Grade (ab74272)

This image is courtesy of an Abreview submitted by Qing Wen

ab74272 staining Androgen Receptor in adult mouse testis tissue sections by Immunohistochemistry (IHC-Fr - frozen sections). Tissue was fixed with 4% PFA and blocked with 5% BSA for 45 minutes at 25°C. Samples were incubated with primary antibody (1/400 in 1% BSA) for 16 hours at 4°C. A TRITC-conjugated donkey anti-rabbit IgG polyclonal (1/400) was used as the secondary antibody.

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors