

Product datasheet

Anti-Axin 2 antibody [EPR2005(2)] ab109307

RabMAb

★★★★☆ 5 Abreviews 3 References 8 图像

概述

产品名称	Anti-Axin 2抗体[EPR2005(2)]
描述	兔单克隆抗体[EPR2005(2)] to Axin 2
宿主	Rabbit
经测试应用	适用于: ICC/IF, IHC-P, WB
种属反应性	与反应: Mouse, Rat, Human, Pig
免疫原	Synthetic peptide (the amino acid sequence is considered to be commercially sensitive) corresponding to Human Axin 2 aa 50-150.
阳性对照	WB: C6, PC-12, NIH/3T3, MCF7, SW480, and PC3 cell lysates. IHC-P: Human lung carcinoma tissue. ICC/IF: LnCap cells.
常规说明	<p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p> <p>This product is a recombinant rabbit monoclonal antibody.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 59% PBS, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR2005(2)

同种型

IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab109307** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ICC/IF	★★★★★	1/150.
IHC-P	★★★★☆	1/150.
WB	★★★★★	1/1000 - 1/2000. Detects a band of approximately 95 kDa (predicted molecular weight: 94 kDa).

靶标


功能	Inhibitor of the Wnt signaling pathway. Down-regulates beta-catenin. Probably facilitate the phosphorylation of beta-catenin and APC by GSK3B.
组织特异性	Expressed in brain and lymphoblast.
疾病相关	Defects in AXIN2 are involved in colorectal cancer (CRC) [MIM:114500]. They appear to be specifically associated with defective mismatch repair. Defects in AXIN2 are the cause of oligodontia-colorectal cancer syndrome (ODCRCS) [MIM:608615]. Affected individuals manifest severe tooth agenesis and colorectal cancer or precancerous lesions of variable types.
序列相似性	Contains 1 DIX domain. Contains 1 RGS domain.
结构域	The tankyrase-binding motif (also named TBD) is required for interaction with tankyrase TNKS and TNKS2.
翻译后修饰	Probably phosphorylated by GSK3B and dephosphorylated by PP2A. ADP-ribosylated by tankyrase TNKS and TNKS2. Poly-ADP-ribosylated protein is recognized by RNF146, followed by ubiquitination and subsequent activation of the Wnt signaling pathway. Ubiquitinated by RNF146 when poly-ADP-ribosylated, leading to its degradation and subsequent activation of the Wnt signaling pathway. Deubiquitinated by USP34, deubiquitinated downstream of beta-catenin stabilization step: deubiquitination is important Wnt signaling to positively regulate beta-catenin (CTNBB1)-mediated transcription.
细胞定位	Cytoplasm.

图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling Axin 2 with unpurified ab109307 at 1/150. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Western blot - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

All lanes : Anti-Axin 2 antibody [EPR2005(2)] (ab109307) at 1/1000 dilution (unpurified)

Lane 1 : SW480 cell lysate

Lane 2 : C6 cell lysate

Lane 3 : PC-12 cell lysate

Lane 4 : NIH/3T3 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 94 kDa

Observed band size: 94 kDa

Blocking buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm /TBST.



Immunocytochemistry/ Immunofluorescence - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)


This image is courtesy of an anonymous Abreview

Unpurified ab109307 staining Axin 2 in 293T cells by ICC/IF

(Immunocytochemistry/immunofluorescence).

Cells were fixed with paraformaldehyde, permeabilized with 1% Triton X-100 and blocked with 3% BSA for 1 hour at room temperature.

Samples were incubated with primary antibody (1/50 in PBS + 3% BSA) for 16 hours. An Alexa Fluor[®] 488-conjugated donkey anti-rabbit IgG polyclonal (1/500) was used as the secondary antibody.



Western blot - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

All lanes : Anti-Axin 2 antibody [EPR2005(2)] (ab109307) at 1/1000 dilution (purified)

Lane 1 : SW480 cell lysate

Lane 2 : C6 cell lysate

Lane 3 : PC-12 cell lysate

Lane 4 : NIH/3T3 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 94 kDa

Observed band size: 94 kDa


Blocking buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling Axin 2 with purified ab109307 at 1/150. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)



Western blot - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

All lanes : Anti-Axin 2 antibody [EPR2005(2)] (ab109307) at 1/1000 dilution (unpurified)

Lane 1 : MCF7 cell lysates


Lane 2 : SW480 cell lysates

Lane 3 : PC3 cell lysates

Lysates/proteins at 10 µg per lane.

Predicted band size: 94 kDa


Observed band size: 95 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

Immunocytochemistry/Immunofluorescence analysis of LnCap cells labelling Axin 2 with unpurified ab109307 at 1/150. Cells were fixed with 4% paraformaldehyde. An Alexa Fluor[®] 555-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/150) and secondary antibody, ab150113, an Alexa Fluor[®] 488-conjugated goat anti-mouse IgG (1/500).



Immunocytochemistry/ Immunofluorescence - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

Immunocytochemistry/Immunofluorescence analysis of LnCap cells labelling Axin 2 with purified ab109307 at 1/150. Cells were fixed with 4% paraformaldehyde. An Alexa Fluor[®] 555-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/150) and secondary antibody, ab150113, an Alexa Fluor[®] 488-conjugated goat anti-mouse IgG (1/500).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors