

Product datasheet

Anti-CD4 antibody [W3/25] (Biotin) ab111815

1 References 1 图像

概述

| | |
|-------|---|
| 产品名称 | Anti-CD4抗体[W3/25] (Biotin) |
| 描述 | 小鼠单克隆抗体[W3/25] to CD4 (Biotin) |
| 宿主 | Mouse |
| 偶联物 | Biotin |
| 经测试应用 | 适用于: Flow Cyt, IHC-Fr, IHC-P |
| 种属反应性 | 与反应: Rat |
| 免疫原 | Rat thymocyte membrane. |
| 阳性对照 | Thymus, Spleen and Lymph Node cells |
| 常规说明 | Abcam is committed to meeting high standards of ethical manufacturing and has decided to discontinue this product by June 2019 as it has been generated by the ascites method. We are sorry for any inconvenience this may cause. |

性能

| | |
|------|---|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| 存储溶液 | Preservative: 0.02% Sodium azide Constituent: 0.5% BSA |
| 纯度 | Protein G purified |
| 克隆 | 单克隆 |
| 克隆编号 | W3/25 |
| 骨髓瘤 | P3-NS1/1-Ag4-1 |
| 同种型 | IgG1 |

应用

Our [Abpromise guarantee](#) covers the use of **ab111815** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| 应用 | Ab评论 | 说明 |
|----------|------|--|
| Flow Cyt | | Use 0.5µg for 10 ⁶ cells. ab18434 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody. |
| IHC-Fr | | Use at an assay dependent concentration. |
| IHC-P | | Use at an assay dependent concentration. |

靶标

| | |
|-------|--|
| 功能 | Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell activation. Induces the aggregation of lipid rafts. |
| 序列相似性 | Contains 3 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 Ig-like V-type (immunoglobulin-like) domain. |
| 翻译后修饰 | Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts. |
| 细胞定位 | Cell membrane. Localizes to lipid rafts. Removed from plasma membrane by HIV-1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation. Cell surface expression is also down-modulated by HIV-1 Envelope polyprotein gp160 that interacts with, and sequesters CD4 in the endoplasmic reticulum. |

图片



[ab11815](#) at 0.5 µg/10⁶ cells staining CD4 in rat strain Buffalo thymus cells by Flow Cytometry.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors