

Product datasheet

Anti-CD40 antibody ab113701

2 图像

概述

产品名称	Anti-CD40抗体
描述	兔多克隆抗体to CD40
宿主	Rabbit
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Human 预测可用于: Macaque monkey, Gorilla 
免疫原	Synthetic peptide conjugated to KLH derived from within residues 50 - 150 of Human CD40. 参阅Abcam的专有抗源政策
阳性对照	This antibody gave a positive signal in Human Spleen and Thymus tissue lysates. This antibody gave a positive result in IHC in the following FFPE tissue: Human normal spleen.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS Note: Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab113701** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 43 kDa (predicted molecular weight: 31 kDa).
IHC-P		Use a concentration of 5 µg/ml.

靶标

功能	Receptor for TNFSF5/CD40LG.
组织特异性	B-cells and in primary carcinomas.
疾病相关	Defects in CD40 are the cause of hyper-IgM immunodeficiency syndrome type 3 (HIGM3) [MIM:606843]; also known as hyper-IgM syndrome 3. HIGM3 is an autosomal recessive disorder which includes an inability of B cells to undergo isotype switching, one of the final differentiation steps in the humoral immune system, an inability to mount an antibody-specific immune response, and a lack of germinal center formation.
序列相似性	Contains 4 TNFR-Cys repeats.
细胞定位	Secreted and Cell membrane.

图片



Western blot - Anti-CD40 antibody (ab113701)

All lanes : Anti-CD40 antibody (ab113701) at 1 µg/ml

Lane 1 : Human spleen tissue lysate - total protein ([ab29699](#))

Lane 2 : Human thymus tissue lysate - total protein ([ab30146](#))

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) preadsorbed ([ab97080](#)) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 31 kDa


Observed band size: 43 kDa

Additional bands at: 105 kDa, 41 kDa

(possible cleavage fragment). We are unsure as to the identity of these extra bands.

Exposure time: 10 minutes

CD40 contains a number of potential glycosylation sites (SwissProt) which may explain its migration at a higher molecular weight than predicted. The predicted molecular weight of CD40 is 31 kDa (SwissProt), however we expect to observe a banding pattern around 43 kDa. Abcam welcomes customer feedback and would appreciate any comments regarding this product and the data presented above.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD40 antibody (ab113701)

IHC image of CD40 staining in Human normal spleen formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab113701, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors