

Product datasheet

Anti-CD40L antibody [5F3] ab13546

概述

| | |
|-------|--|
| 产品名称 | Anti-CD40L抗体[5F3] |
| 描述 | 小鼠单克隆抗体[5F3] to CD40L |
| 宿主 | Mouse |
| 经测试应用 | 适用于: Flow Cyt |
| 种属反应性 | 与反应: Human |
| 免疫原 | Human CD40L (CD154) transfected 300-19 cell (Mouse pre-B cell line). |
| 常规说明 | <p>For maximum product recovery, after thawing, centrifuge the product vial before removing cap.</p> <p>Abcam is committed to meeting high standards of ethical manufacturing and as such, we will be discontinuing this product, which has been generated by the ascites method, within the next year. We are sorry for any inconvenience this may cause. If you would like help finding an alternative product, please do not hesitate to contact our scientific support team.</p> |

性能

| | |
|------|--|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| 存储溶液 | Preservative: None Constituents: 1% Sucrose, PBS, pH 7.2. |
| 纯度 | Protein A purified |
| 纯化说明 | This antibody is affinity purified. |
| 克隆 | 单克隆 |
| 克隆编号 | 5F3 |
| 同种型 | IgG1 |

应用

Our [Abpromise guarantee](#) covers the use of **ab13546** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| 应用 | Ab评论 | 说明 |
|----------|------|---|
| Flow Cyt | | Use a concentration of 5 - 10 µg/ml. ab170190 -Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody. |

靶标

| | |
|--------------|--|
| 功能 | Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL-4. Involved in immunoglobulin class switching. Release of soluble CD40L from platelets is partially regulated by GP IIb/IIIa, actin polymerization, and an matrix metalloproteinases (MMP) inhibitor-sensitive pathway. |
| 组织特异性 | Specifically expressed on activated CD4+ T-lymphocytes. |
| 疾病相关 | Defects in CD40LG are the cause of X-linked immunodeficiency with hyper-IgM type 1 (HIGM1) [MIM:308230]; also known as X-linked hyper IgM syndrome (XHIM). HIGM1 is an immunoglobulin isotype switch defect characterized by elevated concentrations of serum IgM and decreased amounts of all other isotypes. Affected males present at an early age (usually within the first year of life) recurrent bacterial and opportunistic infections, including Pneumocystis carinii pneumonia and intractable diarrhea due to cryptosporidium infection. Despite substitution treatment with intravenous immunoglobulin, the overall prognosis is rather poor, with a death rate of about 10% before adolescence. |
| 序列相似性 | Belongs to the tumor necrosis factor family. |
| 翻译后修饰 | The soluble form derives from the membrane form by proteolytic processing. N-linked glycan is a mixture of high mannose and complex type. Glycan structure does not influence binding affinity to CD40. Not O-glycosylated. |
| 细胞定位 | Secreted and Cell membrane. |

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors