

Product datasheet

Anti-CD44 antibody [EPR1013Y] ab51037

重组 RabMAb


★★★★☆ 10 Abreviews 55 References 11 图像

概述

产品名称	Anti-CD44抗体[EPR1013Y]
描述	兔单克隆抗体[EPR1013Y] to CD44
经测试应用	适用于: IHC-P, IHC-Fr, WB, Flow Cyt 不适用于: IP
种属反应性	与反应: Human 不与反应: Mouse, Rat
免疫原	Synthetic peptide (the amino acid sequence is considered to be commercially sensitive) within Human CD44 aa 150-250. The exact sequence is proprietary.
阳性对照	TF-1 cell lysate or human urinary bladder tissue.
常规说明	Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents This product is a recombinant rabbit monoclonal antibody.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
解离常数 (K_D)	K _D = 3.76 x 10 ⁻¹⁰ M



[Learn more about K_D](#)

存储溶液	PBS 49%, Sodium azide 0.01%, Glycerol 50%, BSA 0.05%
纯度	Tissue culture supernatant
克隆	单克隆
克隆编号	EPR1013Y
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab51037** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
IHC-P	★★★★★	1/50 - 1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
IHC-Fr		Use at an assay dependent concentration.
WB	★★★★☆	1/5000. Detects a band of approximately 82 kDa (predicted molecular weight: 82 kDa).
Flow Cyt	★★★★☆	1/30. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

应用说明 Is unsuitable for IP.

靶标


功能	Receptor for hyaluronic acid (HA). Mediates cell-cell and cell-matrix interactions through its affinity for HA, and possibly also through its affinity for other ligands such as osteopontin, collagens, and matrix metalloproteinases (MMPs). Adhesion with HA plays an important role in cell migration, tumor growth and progression. Also involved in lymphocyte activation, recirculation and homing, and in hematopoiesis. Altered expression or dysfunction causes numerous pathogenic phenotypes. Great protein heterogeneity due to numerous alternative splicing and post-translational modification events.
组织特异性	Isoform 10 (epithelial isoform) is expressed by cells of epithelium and highly expressed by carcinomas. Expression is repressed in neuroblastoma cells.
序列相似性	Contains 1 Link domain.
结构域	The lectin-like LINK domain is responsible for hyaluronan binding.
翻译后修饰	Proteolytically cleaved in the extracellular matrix by specific proteinases (possibly MMPs) in several cell lines and tumors. N-glycosylated. O-glycosylated; contains more-or-less-sulfated chondroitin sulfate glycans, whose number may affect the accessibility of specific proteinases to their cleavage site(s). Phosphorylated; activation of PKC results in the dephosphorylation of Ser-706 (constitutive phosphorylation site), and the phosphorylation of Ser-672.
细胞定位	Membrane.

图片




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)-Anti-CD44 antibody [EPR1013Y](ab51037)

ab51037 (1:100) showing positive staining in human Breast carcinoma tissue.




Flow Cytometry - Anti-CD44 antibody [EPR1013Y] (ab51037)

ab51037 used at a 1/200 dilution (red) to detect CD44 in human MDA-MB-231 cells by Flow Cytometry. The secondary used was a Pgoat anti-rabbit IgG (Alexa Fluor® 488) polyclonal used at a 1/2000 dilution. Isotype control (black): Rabbit monoclonal IgG (ab172730). Unlabelled control (Blue): Cell without incubation with primary antibody and secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)

Immunohistochemical analysis of paraffin-embedded human pancreatic cancer tissue labeling CD44 with ab51037 at 1/100 dilution followed by goat anti-rabbit IgG H&L (HRP) (ab97051, 1/500). The sample was counter stained with hematoxylin.



Western blot - Anti-CD44 antibody [EPR1013Y] (ab51037)

All lanes : Anti-CD44 antibody [EPR1013Y] (ab51037) at 1/1000 dilution

Lane 1 : MCF-7 whole cell lysate

Lane 2 : Jurkat whole cell lysate

Lane 3 : MDA-MB-231 whole cell lysate

Lane 4 : HeLa whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution


Predicted band size : 82 kDa

Observed band size : 81 kDa

Exposure time : 3 minutes


Blocking and diluting buffer 5% NFDM/TBST

The expression of CD44 in MCF-7 is low (PMID: 25635866; PMID: 26005723). Jurkat does not express CD44 (PMID: 24127558).




Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling CD44 with ab51037 at 1/100 dilution followed by goat anti-rabbit IgG H&L (HRP) (ab97051, 1/500). The sample was counter stained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)




Immunohistochemical analysis of paraffin-embedded human skin tissue labeling CD44 with ab51037 at 1/100 dilution followed by goat anti-rabbit IgG H&L (HRP) (ab97051, 1/500). The sample was counter stained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue labeling CD44 with ab51037 at 1/100 dilution followed by goat anti-rabbit IgG H&L (HRP) (ab97051, 1/500). The sample was counter stained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)



Western blot - CD44 antibody [EPR1013Y]
(ab51037)

Anti-CD44 antibody [EPR1013Y] (ab51037)
at 1/5000 dilution + TF-1 cell lysate at 10 μ g

Secondary

Goat anti-Rabbit HRP labeled. at 1/2000
dilution


Predicted band size : 82 kDa

Observed band size : 82 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections)-Anti-CD44 antibody
[EPR1013Y](ab51037)

ab51037 (1:100) showing positive staining in
human Glioma tissue.



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections)-Anti-CD44 antibody
[EPR1013Y](ab51037)

ab51037 (1:100) showing positive staining in
human Thyroid gland carcinoma tissue.



Other-Anti-CD44 antibody [EPR1013Y](ab51037)

Equilibrium disassociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors