

Product datasheet

Anti-Collagen V antibody ab21299

概述

| | |
|-------|--|
| 产品名称 | Anti-Collagen V抗体 |
| 描述 | 兔多克隆抗体to Collagen V |
| 宿主 | Rabbit |
| 特异性 | Cross-Reactivity (RIA%) Rat Collagen Type V 100% Rat Collagen Types I, III <0.1% Rat Elastin <0.1% |
| 经测试应用 | 适用于: ELISA, ICC/IF, RIA |
| 种属反应性 | 与反应: Rat |
| 免疫原 | Collagen type V extracted and purified from rat skin. |

性能

| | |
|------|---|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| 存储溶液 | Buffer: Not applicable. Preservative: None |
| 纯度 | Ion Exchange Chromatography |
| 克隆 | 多克隆 |
| 同种型 | IgG |

应用

Our [Abpromise guarantee](#) covers the use of **ab21299** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| 应用 | Ab评论 | 说明 |
|--------|------|----|
| ELISA | | |
| ICC/IF | | |
| RIA | | |

应用说明 ELISA: Use at >1:100-1:200 dilution.

IF: Use at an assay dependent dilution. In indirect IFA with fluorescein anti-rabbit IgG conjugate, use diluted at >1:40 on frozen rat tissue sections.

(SP)RIA: Use at an assay dependent dilution.

For immunostaining of extra or intracellular components in light microscopy. For the quantitation of Collagen Type V in biological fluids or in culture supernatants.

Each laboratory should determine an optimum working titer for use in its particular application.

靶标

| | |
|-------|---|
| 功能 | Type V collagen is a member of group I collagen (fibrillar forming collagen). It is a minor connective tissue component of nearly ubiquitous distribution. Type V collagen binds to DNA, heparan sulfate, thrombospondin, heparin, and insulin. |
| 疾病相关 | Ehlers-Danlos syndrome 1 Ehlers-Danlos syndrome 2 |
| 序列相似性 | Belongs to the fibrillar collagen family. Contains 1 fibrillar collagen NC1 domain. Contains 1 laminin G-like domain. |
| 结构域 | The C-terminal propeptide, also known as COLFI domain, have crucial roles in tissue growth and repair by controlling both the intracellular assembly of procollagen molecules and the extracellular assembly of collagen fibrils. It binds a calcium ion which is essential for its function. |
| 翻译后修饰 | Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains. Sulfated on 40% of tyrosines. |
| 细胞定位 | Secreted > extracellular space > extracellular matrix. |

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors