

Product datasheet

Anti-Dynorphin A antibody ab82509

★★★★☆ 1 Abreviews 1 References 5 图像

概述

产品名称	Anti-Dynorphin A抗体
描述	兔多克隆抗体to Dynorphin A
宿主	Rabbit
经测试应用	适用于: ICC/IF, IHC-P, WB
种属反应性	与反应: Mouse, Human 预测可用于: Rat, Guinea pig, Cow, Pig, Monkey 
免疫原	Synthetic peptide conjugated to KLH derived from within residues 200 to the C-terminus of Human Dynorphin A.参阅Abcam的专有抗源政策(Peptide available as ab91634 .)
阳性对照	This antibody gave a positive signal in human brain membrane tissue lysate. IHC-P: FFPE mouse dorsal root ganglion.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab82509** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ICC/IF		Use a concentration of 5 µg/ml.

应用	Ab评论	说明
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB	★★★★☆	Use a concentration of 1 µg/ml. Detects a band of approximately 28 kDa (predicted molecular weight: 28 kDa).

靶标


相关性

Dynorphins are a class of peptides produced by many different populations of neurons, which has some opiate like activity. It is thus classed as an endogenous opioid peptide. The dynorphins, which include dynorphin A, dynorphin B, alpha and beta neoendorphin, and big dynorphin, are all the products of a single gene, 'preprodynorphin'. Dynorphin is produced in many different parts of the brain, including the hypothalamus, the hippocampus and the spinal cord, and has many different physiological actions, depending upon its site of production. Dynorphin is involved in modulating responses to several psychoactive substances, including cocaine. As such, it may help some individuals against addiction. Blocking dynorphin may help alleviate depression.

细胞定位


Secreted

图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Dynorphin A antibody (ab82509)

IHC image of Dynorphin A staining in formalin fixed, paraffin embedded mouse dorsal root ganglion tissue section, performed on a Leica Bond™ system using the standard protocol B. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab82509, 10µg/ml, for 15 mins at room temperature. A goat anti-rabbit biotinylated secondary antibody was used to detect the primary, and visualized using an HRP conjugated ABC system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blot - Dynorphin A antibody (ab82509)

Anti-Dynorphin A antibody (ab82509) at 1 $\mu\text{g/ml}$ + Human brain normal tissue lysate - membrane extract (ab29456) at 10 μg

Secondary

Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 28 kDa

Observed band size: 28 kDa


Exposure time: 15 minutes



Immunocytochemistry/ Immunofluorescence - Dynorphin A antibody (ab82509)


ICC/IF image of ab82509 stained PC12 cells.

The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab82509, 5 $\mu\text{g/ml}$) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43 μM . This antibody also gave a positive result in 100% methanol fixed (5 min) PC12 cells at 5 $\mu\text{g/ml}$.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Dynorphin A antibody (ab82509)

IHC image of Dynorphin A staining in human hippocampus formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab82509, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX



Western blot - Anti-Dynorphin A antibody (ab82509)
This image is courtesy of an anonymous Abreview

Anti-Dynorphin A antibody (ab82509) at 1/1000 dilution + Mouse synaptosome from hippocampal stratum lucidum tissue lysate at 10 µg

Secondary

HRP-conjugated anti-rabbit IgG polyclonal at 1/5000 dilution

Developed using the ECL technique.

Performed under non-reducing conditions.

Predicted band size: 28 kDa

Observed band size: 26 kDa

Additional bands at: 56 kDa (possible non-specific binding), 65 kDa (possible non-specific binding)

Exposure time: 15 seconds

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors