


Product datasheet

Anti-EPB41 antibody ab98854

1 图像

概述

产品名称	Anti-EPB41抗体
描述	兔多克隆抗体to EPB41
经测试应用	适用于: WB
种属反应性	与反应: Human 预测可用于: Mouse, Rat, Rabbit, Horse, Chicken, Guinea pig, Cat, Dog 
免疫原	Synthetic peptide corresponding to a region within N terminal amino acids 107 - 156 (EKGEGGQKEI EFGTSLDEEI ILKAPIAAPE PELKTDPSLD LHSLSSAETQ) of Human EPB41 (NP_976218; UniProt ID: P11171 isoform 5). Run BLAST with ExPASy Run BLAST with NCBI
阳性对照	Human fetal spleen lysate

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: None Constituents: 2% Sucrose, PBS
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab98854** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab 评论	说明
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: for isoform 5, 87 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

靶标

功能	Protein 4.1 is a major structural element of the erythrocyte membrane skeleton. It plays a key role in regulating membrane physical properties of mechanical stability and deformability by stabilizing spectrin-actin interaction. Recruits DLG1 to membranes.
疾病相关	<p>Defects in EPB41 are the cause of elliptocytosis type 1 (EL1) [MIM:611804]. EL1 is a Rhesus-linked form of hereditary elliptocytosis, a genetically heterogeneous, autosomal dominant, hematologic disorder. It is characterized by variable hemolytic anemia and elliptical or oval red cell shape.</p> <p>Defects in EPB41 are a cause of hereditary pyropoikilocytosis (HPP) [MIM:266140]. HPP is an autosomal recessive hematologic disorder characterized by hemolytic anemia, microspherocytosis, poikilocytosis, and an unusual thermal sensitivity of red cells.</p>
序列相似性	Contains 1 FERM domain.
翻译后修饰	<p>Phosphorylated at multiple sites by different protein kinases and each phosphorylation event selectively modulates the protein's functions.</p> <p>Phosphorylation on Tyr-660 reduces the ability of 4.1 to promote the assembly of the spectrin/actin/4.1 ternary complex.</p> <p>O-glycosylated; contains N-acetylglucosamine side chains in the C-terminal domain.</p>
细胞定位	Cytoplasm > cytoskeleton. Cytoplasm > cell cortex. Nucleus.

Anti-EPB41 antibody 图像



Anti-EPB41 antibody (ab98854) at 1 µg/ml +
Human fetal spleen lysate at 10 µg

Predicted band size : for isoform 5, 87 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors