

## Product datasheet

# Anti-Eph receptor A8 antibody ab10615

### 概述

<b>产品名称</b>	Anti-Eph receptor A8抗体
<b>描述</b>	山羊多克隆抗体to Eph receptor A8
<b>特异性</b>	Anti-Mouse EphA8 recognizes recombinant mouse EphA8 by immunoblotting and ELISA. The antibody shows approximately 5% cross-reactivity with recombinant rat EphA5, recombinant mouse EphA2, recombinant mouse EphA3, recombinant mouse EphA6, recombinant mouse EphA7, and recombinant human EphA1.
<b>经测试应用</b>	<b>适用于:</b> ELISA, WB
<b>种属反应性</b>	<b>与反应:</b> Mouse
<b>免疫原</b>	Purified recombinant mouse EphA8 extracellular domain.

### 性能

<b>形式</b>	Liquid
<b>存放说明</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>存储溶液</b>	Preservative: None Constituents: PBS
<b>纯度</b>	Immunogen affinity purified
<b>克隆</b>	多克隆
<b>同种型</b>	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab10615** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
----	------	----

ELISA

WB

**应用说明** ELISA: Use at a concentration of 0.5 - 1 µg/ml. For ELISAs, a working antibody concentration of

0.5-1.0 µg/ml is recommended. The detection limit for recombinant mouse EphA8 is approximately 0.3 ng/well.

WB: Use at a concentration of 0.1 - 0.2 µg/ml. Detects a band of approximately 120 kDa (predicted molecular weight: 120 kDa). The detection limit for recombinant mouse EphA8 is approximately 2 ng/lane under non-reducing and reducing conditions.

Not tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

---

## 靶标

功能	Receptor for members of the ephrin-A family.
序列相似性	Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily. Contains 2 fibronectin type-III domains. Contains 1 protein kinase domain. Contains 1 SAM (sterile alpha motif) domain.
翻译后修饰	Phosphorylation on Tyr-616 is critical for association with FYN. Phosphorylation on Tyr-839 modulates tyrosine kinase activity.
细胞定位	Membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

---

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

---

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors