

# FITC Anti-Fibrinogen antibody ab73925

## 1 References

### 概述

|       |   |
|-------|---|
| 产品名称  | FITC荧光Anti-Fibrinogen抗体   |
| 描述    | FITC荧光鸡多克隆抗体to Fibrinogen   |
| 宿主    | Chicken   |
| 偶联物   | FITC. Ex: 493nm, Em: 528nm  |
| 种属反应性 | 与反应: Rat, Rabbit, Human, Pig  |
| 免疫原   | Purified full length human Fibrinogen.  |
| 常规说明  | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p> |

### 性能

|      |   |
|------|---|
| 形式   | Liquid  |
| 存放说明 | Shipped at 4°C. Store at +4°C.  |
| 存储溶液 | pH: 7.20<br>Preservative: 0.1% Sodium azide<br>Constituents: 0.87% Sodium chloride, 0.328% Sodium phosphate         |
| 纯度   | IgY fraction  |
| 纯化说明 | The IgY fraction is isolated by a two step PEG precipitation procedure followed by ammonium sulphate precipitation. |
| 克隆   | 多克隆   |
| 同种型  | IgY   |

### 应用

|      |   |
|------|---|
| 应用说明 | Flow Cyt: Use at 1/10 for the detection of platelet activation. The antibody should be added to |
|------|---|

Platelet-rich plasma prediluted 1/20 with Hepes buffer (pH 7.4). Fix platelets with 1% p-formaldehyde in PBS.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

---

## 靶标

### 功能

Fibrinogen has a double function: yielding monomers that polymerize into fibrin and acting as a cofactor in platelet aggregation.

### 组织特异性

Plasma.

### 疾病相关

Defects in FGA are a cause of congenital afibrinogenemia (CAFBN) [MIM:202400]. This is a rare autosomal recessive disorder characterized by bleeding that varies from mild to severe and by complete absence or extremely low levels of plasma and platelet fibrinogen. Note=The majority of cases of afibrinogenemia are due to truncating mutations. Variations in position Arg-35 (the site of cleavage of fibrinopeptide a by thrombin) leads to alpha-dysfibrinogenemias.

Defects in FGA are a cause of amyloidosis type 8 (AMYL8) [MIM:105200]; also known as systemic non-neuropathic amyloidosis or Ostertag-type amyloidosis. AMYL8 is a hereditary generalized amyloidosis due to deposition of apolipoprotein A1, fibrinogen and lysozyme amyloids. Viscera are particularly affected. There is no involvement of the nervous system. Clinical features include renal amyloidosis resulting in nephrotic syndrome, arterial hypertension, hepatosplenomegaly, cholestasis, petechial skin rash.

### 序列相似性

Contains 1 fibrinogen C-terminal domain.

### 结构域

A long coiled coil structure formed by 3 polypeptide chains connects the central nodule to the C-terminal domains (distal nodules). The long C-terminal ends of the alpha chains fold back, contributing a fourth strand to the coiled coil structure.

### 翻译后修饰

The alpha chain is not glycosylated.

Forms F13A-mediated cross-links between a glutamine and the epsilon-amino group of a lysine residue, forming fibronectin-fibrinogen heteropolymers.

About one-third of the alpha chains in the molecules in blood were found to be phosphorylated.

Conversion of fibrinogen to fibrin is triggered by thrombin, which cleaves fibrinopeptides A and B from alpha and beta chains, and thus exposes the N-terminal polymerization sites responsible for the formation of the soft clot. The soft clot is converted into the hard clot by factor XIIIa which catalyzes the epsilon-(gamma-glutamyl)lysine cross-linking between gamma chains (stronger) and between alpha chains (weaker) of different monomers.

Phosphorylation sites are present in the extracellular medium.

### 细胞定位

Secreted.

---

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

---

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors