

Product datasheet

Anti-GFAP antibody [EPR1034Y] ab68428

RabMAb


★★★★★ 4 Abreviews 10 References 12 图像

概述

产品名称	Anti-GFAP抗体[EPR1034Y]
描述	兔单克隆抗体[EPR1034Y] to GFAP
宿主	Rabbit
经测试应用	适用于: WB, IP, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide within Human GFAP aa 1 to the C-terminus (N terminal). The exact sequence is proprietary.
阳性对照	WB: Human, Mouse and Rat brain tissue lysate; Human, Mouse and Rat cerebellum tissue lysate; IHC-P: Mouse brain, cerebral cortex and liver tissue sections; Human brain, cerebral cortex, hippocampus, colon and glioma tissue sections ICC/IF: Mouse cerebellum IP: Rat brain whole cell lysate
常规说明	<p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p> <p>This product is a recombinant rabbit monoclonal antibody.</p>

性能


形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.21% BSA
纯度	Protein A purified
克隆	单克隆



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [EPR1034Y] (ab68428)


IHC image of GFAP staining in a formalin fixed, paraffin embedded normal human hippocampus tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab68428 at 1/100 dilution for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [EPR1034Y] (ab68428)

Immunohistochemical analysis of formalin-fixed paraffin-embedded human brain (left) and human glioma (right) tissue sections labelling GFAP with unpurified ab68428 at dilution of 1/250.



Western blot - Anti-GFAP antibody [EPR1034Y]
(ab68428)

All lanes : Anti-GFAP antibody [EPR1034Y]
(ab68428) at 1/10000 dilution

Lane 1 : Human cerebellum tissue lysate at
20 µg

Lane 2 : Human brain tissue lysate at 10 µg


Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to
the non-reduced form of IgG at 1/2000 dilution

Predicted band size: 50 kDa


Observed band size: 48-50 kDa

Blocking and Diluting buffer 5% NFDm/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-GFAP antibody
[EPR1034Y] (ab68428)

Immunohistochemical analysis of paraffin-
embedded human cerebral cortex tissue
sections labelling GFAP with purified
ab68428 at a dilution of 1/500. The secondary
antibody used was [ab97051](#), Goat Anti-Rabbit
IgG H&L (HRP) at a dilution of 1/500. The
sample was counterstained with hematoxylin.
Antigen retrieval was performed using EDTA
Buffer; pH 9.0. PBS was used instead of the
primary antibody as the negative control and is
shown in the inset.



Western blot - Anti-GFAP antibody [EPR1034Y]
(ab68428)

All lanes : Anti-GFAP antibody [EPR1034Y]
(ab68428) at 1/10000 dilution

Lane 1 : Mouse cerebellum tissue lysate

Lane 2 : Mouse brain tissue lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to
the non-reduced form of IgG at 1/2000 dilution

Predicted band size: 50 kDa


Observed band size: 50 kDa

Blocking and Diluting buffer 5% NFDm/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-GFAP antibody
[EPR1034Y] (ab68428)

Immunohistochemical analysis of paraffin-
embedded mouse cerebral cortex tissue
sections labelling GFAP with purified
ab68428 at a dilution of 1/500. The secondary
antibody used was [ab97051](#), Goat Anti-Rabbit
IgG H&L (HRP) at a dilution of 1/500. The
sample was counterstained with hematoxylin.
Antigen retrieval was performed using EDTA
Buffer; pH 9.0. PBS was used instead of the
primary antibody as the negative control and is
shown in the inset.



Western blot - Anti-GFAP antibody [EPR1034Y]
(ab68428)

All lanes : Anti-GFAP antibody [EPR1034Y]

(ab68428) at 1/50000 dilution

Lane 1 : Rat cerebellum tissue lysate

Lane 2 : Rat brain tissue lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/2000 dilution

Predicted band size: 50 kDa


Observed band size: 50 kDa

Blocking and Diluting buffer 5% NFDM/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [EPR1034Y] (ab68428)

Immunohistochemical analysis of paraffin-embedded human colon tissue sections labelling GFAP with purified ab68428 at a dilution of 1/500. The secondary antibody used was [ab97051](#), Goat Anti-Rabbit IgG H&L (HRP) at a dilution of 1/500. The sample was counterstained with hematoxylin. Antigen retrieval was performed using EDTA Buffer; pH 9.0.



Immunoprecipitation - Anti-GFAP antibody
[EPR1034Y] (ab68428)

ab68428 at 1/20 dilution
immunoprecipitating GFAP in rat brain whole
cell lysate observed at 50 kDa (lanes 1 and 2).


Lane 1 (input): Rat brain whole cell lysate
10ug

Lane 2 (+): ab68428 + Rat brain whole cell
lysate

Lane 3 (-): Rabbit monoclonal IgG (ab172730)
instead of ab68428 in Rat brain whole cell
lysate

For western blotting, ab68428 was used
followed by VeriBlot for IP (HRP) (ab131366) as
the secondary antibody at a dilution of
1/10,000.

Blocking and Diluting buffer and
concentration: 5% NFDm/TBST.



Western blot - Anti-GFAP antibody [EPR1034Y]
(ab68428)

All lanes : Anti-GFAP antibody [EPR1034Y]
(ab68428) at 1/5000 dilution (unpurified)

Lane 1 : Human brain lysate

Lane 2 : Rat brain lysate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP labelled Goat anti-Rabbit
antibody at 1/2000 dilution


Predicted band size: 50 kDa

Observed band size: 50 kDa



Immunohistochemical analysis of paraffin-embedded mouse liver tissue sections labelling GFAP with purified ab68428 at a dilution of 1/500. The secondary antibody used was ab97051, Goat Anti-Rabbit IgG H&L (HRP) at a dilution of 1/500. The sample was counterstained with hematoxylin. Antigen retrieval was performed using EDTA Buffer; pH 9.0.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [EPR1034Y] (ab68428)



Immunohistochemical analysis of formalin-fixed paraffin-embedded mouse brain tissue section labelling GFAP with unpurified ab68428 at dilution of 1/250.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [EPR1034Y] (ab68428)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors