

## Product datasheet

# Anti-GPCR GPR120 antibody ab118757

2 Abreviews 1 图像

### 概述

产品名称	Anti-GPCR GPR120抗体
描述	兔多克隆抗体to GPCR GPR120
宿主	Rabbit
经测试应用	适用于: WB
种属反应性	与反应: Human 预测可用于: Mouse, Rat, Rabbit, Horse, Guinea pig, Cow, Cat, Dog
免疫原	Synthetic peptide corresponding to a region within internal sequence amino acids 243-292 (VVTHSEITKA SRKRLTVSLA YSESHQIRVS QQDFRLFRTL FLLMVSFFIM) of Human GPCR GPR120. <a href="#">Run BLAST with ExPASy</a> <a href="#">Run BLAST with NCBI</a>
阳性对照	HepG2 cell lysate.

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	Constituents: 97% PBS, 2% Sucrose
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab118757** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
----	------	----

应用	Ab评论	说明
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 42 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

## 靶标

<b>功能</b>	Receptor for medium and long-chain free fatty acid (FAA). Signals via a G(q)/G(11)-coupled pathway. Acts as a receptor for omega-3 fatty acids and mediates robust anti-inflammatory effects particularly in macrophages and fat cells. The anti-inflammatory effects involve inhibition of TAK1 through a beta-arrestin 2 (ARRB2)/TAB1 dependent effect but independent of G(q)/G(11)-coupled pathway. Mediates potent insulin sensitizing and antidiabetic effects by repressing macrophage-induced tissue inflammation. May mediate the taste of fatty acids.
<b>组织特异性</b>	Abundant expression in the intestinal tract.
<b>序列相似性</b>	Belongs to the G-protein coupled receptor 1 family.
<b>翻译后修饰</b>	Phosphorylated. FFA stimulation facilitates phosphorylation.
<b>细胞定位</b>	Cell membrane. Colocalized with ARRB2 following DHA treatment.

## 图片



Anti-GPCR GPR120 antibody (ab118757) at 1 µg/ml + HepG2 cell lysate at 10 µg

**Predicted band size: 42 kDa**

Gel concentration: 12%

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors