

Anti-GSTM1 + GSTM2 antibody ab27489

[3 References](#) [1 图像](#)

概述

产品名称	Anti-GSTM1 + GSTM2抗体
描述	山羊多克隆抗体to GSTM1 + GSTM2
宿主	Goat
特异性	This antibody is expected to recognise both reported variants of human GSTM1 protein (NP_000552; NP_666533) and GSTM2 protein (NP_000848).
经测试应用	适用于: ELISA, WB
种属反应性	与反应: Human 预测可用于: Chimpanzee, Cynomolgus monkey 
免疫原	Synthetic peptide: CVFSKMAVWGK from the C Terminus of the protein (Human). Run BLAST with Run BLAST with
阳性对照	Human liver and lung lysates for Western blot.

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 0.5% Tris, 0.5% BSA
纯度	Immunogen affinity purified
纯化说明	This antibody purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab27489于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
ELISA		1/1000.
WB		Use a concentration of 1 - 3 µg/ml. Detects a band of approximately 26 kDa (predicted molecular weight: 26 kDa). 1 hour primary incubation is recommended for this product.

靶标

相关性 Glutathione S transferases are encoded by two distinct supergene families and a total of eight distinct classes of the soluble cytoplasmic mammalian glutathione S transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. The mu class of these enzymes, to which GSTM1 and 2 belong, functions in the detoxification of electrophilic compounds by conjugation with glutathione. The genes encoding the mu class of enzymes are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs and null mutations of this gene have been linked with an increase in a number of cancers.

细胞定位 Cytoplasmic

图片



Anti-GSTM1 + GSTM2 antibody (ab27489) at 0.3 µg/ml + human lung lysate (35µg protein in RIPA buffer).

Predicted band size: 26 kDa

Observed band size: 26 kDa

Primary incubation was 1 hour. Detected by chemiluminescence.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors