

Product datasheet

Goat Anti-Mouse IgG H&L (TRITC) ab6786

10 References 1 图像

概述

产品名称	山羊Anti-Mouse IgG H&L (TRITC)
靶标种属	Mouse
经测试应用	适用于: Immunomicroscopy, Flow Cyt, IHC-P, IHC-Fr, ICC/IF, ELISA
免疫原	Mouse IgG whole molecule
偶联物	TRITC. Ex: 547nm, Em: 572nm

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 0.878% Sodium chloride, 1% BSA, 0.424% Potassium phosphate
纯度	IgG fraction
纯化说明	This product was prepared from monospecific antiserum by immunoaffinity chromatography using mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
共轭说明	Tetramethylrhodamine isothiocyanate (TRITC) (Molecular Weight 444 daltons) Absorption Wavelength: 550 nm Emission Wavelength: 570 nm Fluorochrome/Protein Ratio: 2.8 moles TRITC per mole of Rabbit IgG
克隆	多克隆
同种型	IgG

应用


Our [Abpromise guarantee](#) covers the use of **ab6786** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
Immunomicroscopy		Use at an assay dependent concentration.

应用	Ab评论	说明
Flow Cyt		1/2000 - 1/4000.
IHC-P		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.
ICC/IF		1/1000 - 1/5000. (PubMed:15659422)
ELISA		1/10000 - 1/50000.

图片



Overlay histogram showing Jurkat cells stained with [ab8090](#) (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody ([ab8090](#), 0.1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody Goat anti-mouse IgG H&L (TRITC) ([ab6786](#)) was used at 1/4000 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIGG2A] ([ab91361](#), 0.1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 561nm laser and 610/20 bandpass filter.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors