

Product datasheet

Anti-HDAC1 antibody ab53091

★★★★★ 6 Abreviews 2 References 8 图像

概述

产品名称	Anti-HDAC1抗体
描述	兔多克隆抗体to HDAC1
宿主	Rabbit
特异性	ab53091 detects endogenous levels of total HDAC1 protein.
经测试应用	适用于: Flow Cyt, ICC, WB, IHC-P, ELISA, ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide derived from human HDAC1
阳性对照	WB: NIH/3T3 cell extracts IHC-P: Human lung carcinoma tissue

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS, 150mM Sodium chloride, pH 7.4
纯度	Immunogen affinity purified
纯化说明	ab53091 was affinity purified from rabbit antiserum by affinity chromatography using epitope specific immunogen.
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab53091** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
Flow Cyt	★★★★★	Use at an assay dependent concentration. ab171870 -Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.

应用	Ab评论	说明
ICC	★★★★★	Use at an assay dependent concentration.
WB	★★★★★	1/300 - 1/1000. Detects a band of approximately 60 kDa (predicted molecular weight: 55 kDa).
IHC-P		Use at an assay dependent concentration.
ELISA		1/10000.
ICC/IF	★★★★★	Use a concentration of 1 µg/ml.

靶标


功能	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Deacetylates SP proteins, SP1 and SP3, and regulates their function. Component of the BRG1-RB1-HDAC1 complex, which negatively regulates the CREST-mediated transcription in resting neurons. Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation. Deacetylates TSHZ3 and regulates its transcriptional repressor activity. Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B.
组织特异性	Ubiquitous, with higher levels in heart, pancreas and testis, and lower levels in kidney and brain.
序列相似性	Belongs to the histone deacetylase family. HD type 1 subfamily.
翻译后修饰	Sumoylated on Lys-444 and Lys-476; which promotes enzymatic activity. Desumoylated by SENP1. Phosphorylation on Ser-421 and Ser-423 promotes enzymatic activity and interactions with NuRD and SIN3 complexes. Ubiquitinated by CHFR, leading to its degradation by the proteasome.
细胞定位	Nucleus.

图片



Immunocytochemistry/ Immunofluorescence-HDAC1 antibody(ab53091)


ICC/IF image of ab53091 stained HeLa cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab53091, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Peptide - +

Immunohistochemistry (Paraffin-embedded sections) - HDAC1 antibody (ab53091)


Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using ab53091 at 1/50 dilution, in the presence (right) or absence (left) of blocking peptide.



Immunocytochemistry/ Immunofluorescence - Anti-HDAC1 antibody (ab53091)

This image is courtesy of an abreview submitted by Dr Mahesh Shivananjappa.

Immunocytochemistry/ Immunofluorescence analysis of mouse RAW 264.7 cells labeling HDAC1 with ab53091 at 1/150 dilution. The cells were fixed with paraformaldehyde and permeabilized with 0.1% Triton-X100 in 2% BSA for 15min. Blocking of the cells was done with 2% BSA for 1 hour at 4°C, followed by staining with ab53091 at 1/150 in 2% BSA in PBS for 16 hours at 4°C. An Alexa Fluor® 488 conjugated chicken anti-rabbit IgG (H+L) secondary antibody was used at 1/1000 dilution.




All lanes : Anti-HDAC1 antibody (ab53091)
at 1/300 dilution

Lane 1 : NIH/3T3 cell extracts

Lane 2 : NIH/3T3 cell extracts with blocking peptide

Predicted band size: 55 kDa


Observed band size: 60 kDa



Flow Cytometry analysis of human mononuclear cell preparations labeling HDAC1 with ab53091 at 1/100 dilution (green). Cells were harvested using a apheresis preparation method. Cells were then fixed with paraformaldehyde and permeabilized with 0.1% Triton-X100 in 2% BSA for 15 min. An Alexa Fluor[®] 488 conjugated chicken anti-rabbit IgG (H+L) was used as the secondary antibody at 1/500 dilution. Blue - secondary only, red - unstained cells.

Flow Cytometry - Anti-HDAC1 antibody (ab53091)

This image is courtesy of an abreview submitted by Dr. Mahesh Shivananjappa.




All lanes : Anti-HDAC1 antibody (ab53091)

Lane 1 : NIH/3T3 cell lysates

Lane 2 : NIH/3T3 cell lysates blocked with synthesized peptide


Predicted band size: 55 kDa

Western blot - Anti-HDAC1 antibody (ab53091)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labeling HDAC1 with ab53091. The image on the right is blocked with the synthesized peptide.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC1 antibody (ab53091)



Immunocytochemistry/ Immunofluorescence analysis of NIH/3T3 cells labeling HDAC1 with ab53091. The image on the right is blocked with the synthesized peptide.

Immunocytochemistry/ Immunofluorescence - Anti-HDAC1 antibody (ab53091)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors