

Product datasheet

Anti-Histone H1.5 antibody ab24175

1 Abreviews 12 References 2 图像

概述

产品名称	Anti-Histone H1.5抗体
描述	兔多克隆抗体to Histone H1.5
宿主	Rabbit
特异性	This antibody is specific for Histone H1.5. Reacts with histone H1 acid-extracted from cell lines such as breast cancer cell line T47D. Also recognizes H1 in a nuclear extract or chromatin preparation. H1.5 is present in many human cell lines.
经测试应用	适用于: ICC/IF, WB, IP
种属反应性	与反应: Human
免疫原	Synthetic peptide: AETATPAPVEKSPAKKKATKKAAGAGAAK -C , corresponding to N terminal amino acids 7-35 of Human Histone H1.5 Run BLAST with Run BLAST with
阳性对照	This antibody gave a positive result in IF/ICC when used in the following formaldehyde fixed cell lines: HepG2.
常规说明	ab24175 does not appear to work in IF in human or mouse cell lines.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Add glycerol to a final volume of 50% for extra stability and aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Constituents: 8% Glycerol
纯度	Protein A purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab24175** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ICC/IF		1/50. PubMed: 19336418
WB		1/500 - 1/1000. Detects a band of approximately 33 kDa (predicted molecular weight: 22 kDa).
IP		Use at an assay dependent dilution. PubMed: 21383955

靶标

功能	Histones H1 are necessary for the condensation of nucleosome chains into higher order structures.
序列相似性	Belongs to the histone H1/H5 family. Contains 1 H15 (linker histone H1/H5 globular) domain.
细胞定位	Nucleus. Chromosome.

图片



All lanes : Histone H1.5 antibody (ab24175)

Lane 1 : Chromatin preparation from a breast cancer cell line

Lane 2 : recombinant human H1.0

Lane 3 : recombinant human H1.1

Lane 4 : recombinant human H1.2

Lane 5 : recombinant human H1.3

Lane 6 : recombinant human H1.4

Lane 7 : recombinant human H1.5

Panel B is a silver stain of the gel in panel A.

All lanes : Histone H1.5 antibody (ab24175)

Lane 1 : Chromatin preparation from a breast cancer cell line

Lane 2 : recombinant human H1.0

Lane 3 : recombinant human H1.1


Lane 4 : recombinant human H1.2

Lane 5 : recombinant human H1.3

Lane 6 : recombinant human H1.4

Lane 7 : recombinant human H1.5

Panel B is a silver stain of the gel in panel A.



Immunocytochemistry/ Immunofluorescence - Anti-Histone H1.5 antibody (ab24175)

ICC/IF image of ab24175 stained HepG2 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab24175 at 5µg/ml overnight at +4°C. The secondary antibody (green) was DyLight® 488 goat anti- rabbit ([ab96899](#)) IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors