

Product datasheet

Anti-Histone H4 (di methyl K79) antibody ab2885

★★★★☆ 2 Abreviews 1 References 4 图像

概述

产品名称	Anti-Histone H4 (di methyl K79)抗体
描述	兔多克隆抗体to Histone H4 (di methyl K79)
宿主	Rabbit
特异性	Specific to Histone H4 K79 Di-methyl. In blocking experiments, is not blocked by H3 K79 peptides.
经测试应用	适用于: WB, IHC-P, ICC/IF, ICC
种属反应性	与反应: Cow, Human 不与反应: Saccharomyces cerevisiae
免疫原	Synthetic peptide within Human Histone H4 aa 50 to the C-terminus (di methyl K79) conjugated to Keyhole Limpet Haemocyanin (KLH). The exact sequence is proprietary. (Peptide available as ab4560)
阳性对照	Calf Thymus Histone Preparation; Hela whole cell extract

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab2885** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB	★★★★☆	1/500. Detects a band of approximately 14 kDa.
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
ICC/IF		1/200.
ICC		1/200.

靶标

功能

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

序列相似性

Belongs to the histone H4 family.

翻译后修饰

Acetylation at Lys-6 (H4K5ac), Lys-9 (H4K8ac), Lys-13 (H4K12ac) and Lys-17 (H4K16ac) occurs in coding regions of the genome but not in heterochromatin.

Citrullination at Arg-4 (H4R3ci) by PADI4 impairs methylation.

Monomethylation and asymmetric dimethylation at Arg-4 (H4R3me1 and H4R3me2a, respectively) by PRMT1 favors acetylation at Lys-9 (H4K8ac) and Lys-13 (H4K12ac). Demethylation is performed by JMJD6. Symmetric dimethylation on Arg-4 (H4R3me2s) by the PRDM1/PRMT5 complex may play a crucial role in the germ-cell lineage.

Monomethylated, dimethylated or trimethylated at Lys-21 (H4K20me1, H4K20me2, H4K20me3). Monomethylation is performed by SET8. Trimethylation is performed by SUV420H1 and SUV420H2 and induces gene silencing.

Phosphorylated by PAK2 at Ser-48 (H4S47ph). This phosphorylation increases the association of H3.3-H4 with the histone chaperone HIRA, thus promoting nucleosome assembly of H3.3-H4 and inhibiting nucleosome assembly of H3.1-H4.

Ubiquitinated by the CUL4-DDB-RBX1 complex in response to ultraviolet irradiation. This may weaken the interaction between histones and DNA and facilitate DNA accessibility to repair proteins. Monoubiquitinated at Lys-92 of histone H4 (H4K91ub1) in response to DNA damage. The exact role of H4K91ub1 in DNA damage response is still unclear but it may function as a licensing signal for additional histone H4 post-translational modifications such as H4 Lys-21 methylation (H4K20me).


Sumoylated, which is associated with transcriptional repression.

Crotonylation (Kcr) is specifically present in male germ cells and marks testis-specific genes in post-meiotic cells, including X-linked genes that escape sex chromosome inactivation in haploid cells. Crotonylation marks active promoters and enhancers and confers resistance to transcriptional repressors. It is also associated with post-meiotically activated genes on autosomes.

细胞定位

Nucleus. Chromosome.

图片



Western blot - Anti-Histone H4 (di methyl K79) antibody (ab2885)

All lanes : Anti-Histone H4 (di methyl K79) antibody (ab2885) at 1/500 dilution

Lane 1 : Calf Thymus histone prep

Lane 2 : Calf Thymus histone prep with Human Histone H3 (mono methyl K79) peptide (ab4555) at 0.1 µg

Lane 3 : Calf Thymus histone prep with Human Histone H3 (di methyl K79) peptide (ab4556) at 0.1 µg

Lane 4 : Calf Thymus histone prep with Human Histone H3 (tri methyl K79) peptide (ab4557) at 0.1 µg

Lane 5 : Calf Thymus histone prep with Human Histone H4 (di methyl K79) peptide (ab4560) at 0.1 µg

Lane 6 : Calf Thymus histone prep with Human Histone H3 (di methyl K27) peptide (ab1781) at 0.1 µg

Lysates/proteins at 1 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab6721) at 1/2000 dilution

Rabbit polyclonal to Histone H4 Di Methyl K79 at 1/500 on Calf Thymus histone prep (1 µg per lane).

Lane 1: ab2885 1/500

Lane 2: ab2885 1/500 H3 Mono Me K79 peptide (ab4555), 0.1 µg


Lane 3: ab2885 1/500 H3 Di Me K79 peptide (ab4556), 0.1 µg

Lane 4: ab2885 1/500 H3 Tri Me K79 peptide (ab4557), 0.1 µg

Lane 5: ab2885 1/500 H4 Di Me K79 peptide (ab4560), 0.1 µg

Lane 6: ab2885 1/500 H3 Di Me K27 peptide (ab1781), 0.1 µg


Secondary antibody: Goat anti-rabbit IgG (HRP) ab 1/2000 (ab6721)



Immunocytochemistry/ Immunofluorescence - Anti-Histone H4 (di methyl K79) antibody (ab2885)

This image is courtesy of Darin McDonal, Hendzel Laboratory

SKN cells were stained with ab2885 (green) at a dilution of 1/200. The cells were fixed in paraformaldehyde for 10 minutes prior to incubation with ab2885. The DNA was stained with DAPI (blue). 100x magnification.



Western blot - Anti-Histone H4 (di methyl K79) antibody (ab2885)

This image is courtesy of an Abreview submitted by Dr Gerald Davies


ab2885 at 1/2000 used in Western blot analysis of MCF7 cells (human breast cancer cells).

Lane 1: MCF7 cell lysate control

Lane 2: MCF7 cell lysate (cells treated with 50 μ m Troglitazone for 48 hours)

Lane 3: MCF7 cells lysate (cells treated with 1 μ m Trichostatin for 48 hours)

Troglitazone and trichostatin increase dimethyl H4 (K79) and act as anti-proliferatives in human breast cancer. The blot was stripped and reprobbed with an alpha tubulin antibody to show loading equivalence.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H4 (di methyl K79) antibody (ab2885)

IHC image of Histone H4 (di methyl K79) staining in human breast carcinoma FFPE section, performed on a Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab2885, 5µg/ml, for 8 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors