

## Product datasheet

### Human EIF2AK1 peptide ab30694

#### 概述

---

产品名称 人EIF2AK1多肽

#### 描述

---

性质 Synthetic

#### 氨基酸序列

种属 Human

#### 技术指标

---

Our [Abpromise guarantee](#) covers the use of **ab30694** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

形式 Liquid

#### 补充说明

- *First try to dissolve a small amount of peptide in either water or buffer. The more charged residues on a peptide, the more soluble it is in aqueous solutions.*
- *If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or buffer.*
- *Consider that any solvent used must be compatible with your assay. If a peptide does not dissolve and you need to recover it, lyophilise to remove the solvent.*
- *Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is cloudy or has gelled the peptide may be in suspension rather than solubilised.*
- *Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior to use.*

#### 制备和贮存

---

稳定性和存储 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Information available upon request.

#### 常规信息

---

<b>功能</b>	Inhibits protein synthesis at the translation initiation level, in response to various stress conditions, including oxidative stress, heme deficiency, osmotic shock and heat shock. Exerts its function through the phosphorylation of EIF2S1 at 'Ser-48' and 'Ser-51', thus preventing its recycling. Binds heme forming a 1:1 complex through a cysteine thiolate and histidine nitrogenous coordination. This binding occurs with moderate affinity, allowing it to sense the heme concentration within the cell. Thanks to this unique heme-sensing capacity, plays a crucial role to shut off protein synthesis during acute heme-deficient conditions. In red blood cells (RBCs), controls hemoglobin synthesis ensuring a coordinated regulation of the synthesis of its heme and globin moieties. Thus plays an essential protective role for RBC survival in anemias of iron deficiency. Similarly, in hepatocytes, involved in heme-mediated translational control of CYP2B and CYP3A and possibly other hepatic P450 cytochromes. May also contain ER stress during acute heme-deficient conditions.
<b>组织特异性</b>	Expressed predominantly in erythroid cells. At much lower levels, expressed in hepatocytes (at protein level).
<b>序列相似性</b>	Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. GCN2 subfamily. Contains 2 HRM (heme regulatory motif) repeats. Contains 1 protein kinase domain.
<b>翻译后修饰</b>	Activated by autophosphorylation; phosphorylated predominantly on serine and threonine residues, but also on tyrosine residues. Autophosphorylation at Thr-488 is required for kinase activation. The active autophosphorylated form apparently is largely refractory to cellular heme fluctuations.
<b>细胞定位</b>	Cytoplasm.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors