

Product datasheet

Human RNF8 peptide ab23278

概述

产品名称 人RNF8多肽

描述

性质 Synthetic

氨基酸序列

种属 Human

序列 GEPGFFVTGDRAG-C

氨基酸 2 to 14

技术指标

Our [Abpromise guarantee](#) covers the use of **ab23278** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用 Blocking - Blocking peptide for Anti-RNF8 antibody ([ab15850](#))

形式 Liquid

制备和贮存

稳定性和存储 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

常规信息

功能 E3 ubiquitin-protein ligase required for assembly of repair proteins to sites of DNA damage. Catalyzes the 'Lys-63'-linked ubiquitination of histone H2A and H2AX. Following DNA double-strand breaks (DSBs), it is recruited to the sites of damage by ATM-phosphorylated MDC1, mediates the ubiquitination of histones H2A and H2AX, thereby promoting the formation of TP53BP1 and BRCA1 ionizing radiation-induced foci (IRIF). Promotes the formation of 'Lys-63'-linked polyubiquitin chains and functions with the specific ubiquitin-conjugating UBE2N/UBC13. Substrates that are polyubiquitinated at 'Lys-63' are usually not targeted for degradation. Enforces the G2/M DNA damage checkpoint. Controls the recruitment of UIMC1-BRCC3

(RAP80-BRCC36) and PAXIP1/PTIP to DNA damage sites following DNA double-strand breaks (DSBs). Ubiquitination of histone H2A requires UBE2N but not MMS2 (UBE2V2). May also ubiquitinate histone H2B. Catalyzes the 'Lys-63'-linked ubiquitination of PCNA. May be required for proper exit from mitosis after spindle checkpoint activation and may regulate cytokinesis. May play a role in the regulation of RXRA-mediated transcriptional activity. Not involved in RXRA ubiquitination by UBE2E2.

组织特异性	Ubiquitous. In fetal tissues, highest expression in brain, thymus and liver. In adult tissues, highest levels in brain and testis, lowest levels in peripheral blood cells.
通路	Protein modification; protein ubiquitination.
序列相似性	Belongs to the RNF8 family. Contains 1 FHA domain. Contains 1 RING-type zinc finger.
发展阶段	Low levels at the G1-S boundary increase in intensity during S phase and until the end of the G2 phase. Abruptly decreases in late mitosis (at protein level). Barely detectable in anaphase.
结构域	The FHA domain specifically recognizes and binds ATM-phosphorylated MDC1 and Thr-4827 phosphorylated HERC2.
翻译后修饰	Autoubiquitinated through 'Lys-48' and 'Lys-63' of ubiquitin. 'Lys-63' polyubiquitination is mediated by UBE2N. 'Lys-29'-type polyubiquitination is also observed, but it doesn't require its own functional RING-type zinc finger.
细胞定位	Nucleus. Midbody. Following DNA double-strand breaks, recruited to the sites of damage. During prophase, concomitant with nuclear envelope breakdown, localizes throughout the cell, with a dotted pattern. In telophase, again in the nucleus and also with a discrete dotted pattern in the cytoplasm. In late telophase and during cytokinesis, localizes in the midbody of the tubulin bridge joining the daughter cells. Does not seem to be associated with condensed chromosomes at any time during the cell cycle.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors