

## Product datasheet

# Anti-MMP9 antibody ab38898

★★★★☆ 43 Abreviews 240 References 8 图像

### 概述

产品名称	Anti-MMP9抗体
描述	兔多克隆抗体to MMP9
宿主	Rabbit
特异性	The antibody binds to Gelatinase-B, but does not cross react with the other MMP family members (MMP-1, MMP-2, MMP-3). In our hands, we observe a weaker signal in WB in human samples compared to mouse samples (BLAST of full length mouse protein sequence showed 72% homology with the Human MMP9 sequence).
经测试应用	适用于: IHC-P, IHC-Fr, WB, IP, ELISA, ICC/IF, ICC, IHC-FoFr
种属反应性	与反应: Mouse, Rat, Dog, Human
免疫原	Full length protein corresponding to Mouse MMP9.
阳性对照	HL60 cell lysate. U937 cell lysate. HT1080 cell lysate. Raw 264.7 cell lysate (LPS treated)

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C.
存储溶液	Preservative: 0.05% Sodium Azide Constituents: 50% Glycerol
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab38898** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
IHC-P	★★★★☆	1/100 - 1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

应用	Ab评论	说明
IHC-Fr	★★★★☆	1/1000. (see Abreview submitted by Greg Gibson) We recommend using <a href="#">Goat Anti-Rabbit IgG H&amp;L (Cy3®) preadsorbed (ab6939) secondary antibody</a>
WB	★★★★☆	1/1000. Detects a band of approximately 92 kDa. When using colorimetric substrates such as BCIP/NBT use at a 1/5000 dilution (for chemiluminescent substrates). Detects a band of approximately 92-95 kDa (pro-form) and 82kDa (active form) (Human samples). Mouse MMP9 is larger, and on SDS PAGE gels runs about 102-105 kDa. Dilution optimised using Chromogenic detection.
IP		Use at an assay dependent concentration.
ELISA	★★★★☆	Use at an assay dependent concentration.
ICC/IF	★★★★☆	1/500.
ICC		Use at an assay dependent concentration.
IHC-FoFr	★★★★★	Use at an assay dependent concentration. PubMed: 19295156

## 靶标

<b>功能</b>	May play an essential role in local proteolysis of the extracellular matrix and in leukocyte migration. Could play a role in bone osteoclastic resorption. Cleaves KiSS1 at a Gly-Leu bond. Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N-terminal one quarter fragments. Degrades fibronectin but not laminin or Pz-peptide.
<b>组织特异性</b>	Produced by normal alveolar macrophages and granulocytes.
<b>疾病相关</b>	Intervertebral disc disease Metaphyseal anadysplasia 2
<b>序列相似性</b>	Belongs to the peptidase M10A family. Contains 3 fibronectin type-II domains. Contains 4 hemopexin repeats.
<b>结构域</b>	The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.
<b>翻译后修饰</b>	Processing of the precursor yields different active forms of 64, 67 and 82 kDa. Sequentially processing by MMP3 yields the 82 kDa matrix metalloproteinase-9. N- and O-glycosylated.
<b>细胞定位</b>	Secreted, extracellular space, extracellular matrix.

## 图片



Western blot - Anti-MMP9 antibody (ab38898)

**All lanes** : Anti-MMP9 antibody (ab38898) at 2 µg/ml

**Lane 1** : Natural human MMP9 protein (Proenzyme, monomer) (ab157344)

**Lane 2** : Recombinant Mouse MMP9 protein (ab39309)


Lysates/proteins at 0.1 µg per lane.

### Secondary

**All lanes** : Infrared labelled goat anti-rabbit (green) at 1/20000 dilution

Performed under reducing conditions.

This blot was produced using a 4-12% Bis-Tris gel under the MOPS buffer system. The gel was run at 200V for 60 minutes before being transferred onto a nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour before being incubated with anti-MMP9 antibody (ab38898; 2 microgram per mL) overnight at 4°C. Antibody binding was detected using infrared labelled goat anti-rabbit (green) antibody (diluted 1:20000) for 1 hour at room temperature before imaging.




Western blot - Anti-MMP9 antibody (ab38898)

Anti-MMP9 antibody (ab38898) + Human  
MMP9

**Observed band size:** 88,92 kDa

Ab38898 detects a band at 92 Kd (pro-form) and a band at 88 Kd (active form). Mouse MMP9 is slightly larger than human MMP9, and the antibody detects a band at about 105 Kd. It is recommend to concentrate samples by Gelatin-agarose affinity chromatography prior to Western blot usage. A recommended starting concentration for Western blots is 1:1000 when using colorimetric substrates such as BCIP/NBT, and 1:5000 for chemiluminescent substrates. Higher concentration of antibody may be needed for non-human samples.



**All lanes** : Anti-MMP9 antibody (ab38898) at 5 µg/ml

**Lane 1** : Recombinant Human MMP9, His tagged ([ab82955](#)) at 0.1 µg

**Lane 2** : U937 whole cell lysate at 100 µg

**Lane 3** : U937 whole cell lysate - treated with PMA and Brefeldin (24 hour treatment) at 100 µg


**Lane 4** : Raw 264.7 (Mouse) whole cell lysate at 100 µg

**Lane 5** : Raw 264.7 (Mouse) whole cell lysate - treated with LPS (6 hour treatment, 1ug/mL) at 100 µg

Performed under reducing conditions.

ab38898 detects recombinant Human MMP9 running at ~85 kDa, and endogenous full-length MMP9 in LPS-stimulated cells at ~100 kDa. This antibody also detects a band at 90 kDa in U937 PMA-treated cells.

ab38898 was incubated at 5 ug/mL and [ab8245](#) (loading control to GAPDH) was diluted at 0.1 ug/mL and both were incubated for 48 hours at 4°C. Blots were developed with goat anti-rabbit IgG (H + L) and goat anti-mouse IgG (H + L) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



**All lanes** : Anti-MMP9 antibody (ab38898) at 1/1000 dilution

**Lanes 1-2** : Human lung tissue lysate at 100 µg with 10% Milk for 1 hour at room temperature

**Lanes 3-4** : MMP-9 KO mice tissue lysates with 10% Milk for 1 hour at room temperature


### Secondary

**All lanes** : HRP-conjugated donkey anti-rabbit polyclonal at 1/1000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Specific observed bands 95-100 kDa



Immunocytochemistry/ Immunofluorescence - Anti-MMP9 antibody (ab38898)

Image courtesy of an anonymous Abreview


ab38898 staining MMP9 (red) in Mouse Neutrophils and Monocytes cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde and permeabilized with 2%BSA + 0.2% tritonX100 in PBS. Samples were incubated with primary antibody (1/200 in 2%BSA + 0.2% tritonX100 in PBS) for 25 minutes at 23°C. An Alexa Fluor® 568-conjugated Donkey anti-rabbit IgG polyclonal (1/1000) was used as the secondary antibody. DAPI is stained blue



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MMP9 antibody (ab38898)

This image is courtesy of an anonymous Abreview

ab38898 staining MMP9 in Mouse Pancreatic carcinoma tissue sections by IHC-P (formaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 1% BSA for 1 hour at room temperature. Antigen retrieval was by heat mediation in citric acid (pH6). Samples were incubated with primary antibody (1/100) in 1% Aurion BSA for 12 hours. An HRP-conjugated Donkey polyclonal to rabbit IgG (1/100) was used as secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MMP9 antibody (ab38898)

Image from Jung IH et al., PLoS One. 2011;6(12):e27941. Epub 2011 Dec 2. Fig 7.; doi:10.1371/journal.pone.0027941; December 2, 2011, PLoS ONE 6(12): e27941.

ab38898 staining MMP9 in 6 month-old transgenic zebrafish pancreas (Ihha overexpression) by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections).

Sections were incubated with primary antibody (1/500) and HRP-conjugated secondary antibody colored using DAB solution. Slides were counterstained with hematoxylin.



Immunohistochemistry (Frozen sections) - Anti-MMP9 antibody (ab38898)

ab38898 at a 1/1000 dilution staining mouse heart tissue by Immunohistochemistry (Frozen sections). The tissue was removed from a mouse, rinsed in PBS and slowly frozen in supercooled isopentane. 14um sections were made using a cryostat. The sections were acetone fixed and blocked in 2% BSA prior to incubation with the MMP9 antibody. [Goat Anti-Rabbit IgG H&L \(Cy3®\) preadsorbed \(ab6939\)](#) was used as the secondary antibody. In the image: red staining = MMP9, blue staining = nuclei, green = gelatinase activity.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors