

Product datasheet

Anti-MMP9 antibody ab38898

★★★★☆ 42 Abreviews 190 References 8 图像

概述

产品名称	Anti-MMP9抗体
描述	兔多克隆抗体to MMP9
特异性	The antibody binds to Gelatinase-B, but does not cross react with the other MMP family members (MMP-1, MMP-2, MMP-3). In our hands, we observe a weaker signal in WB in human samples compared to mouse samples (BLAST of full length mouse protein sequence showed 72% homology with the Human MMP9 sequence).
经测试应用	适用于: IHC-P, IHC-Fr, WB, IP, ELISA, ICC/IF, ICC, IHC-FoFr
种属反应性	与反应: Mouse, Rat, Dog, Human
免疫原	Full length protein corresponding to Mouse MMP9.
阳性对照	HL60 cell lysate. U937 cell lysate. HT1080 cell lysate. Raw 264.7 cell lysate (LPS treated)

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C.
存储溶液	Preservative: 0.05% Sodium Azide Constituents: 50% Glycerol
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab38898** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
IHC-P	★★★★☆	1/100 - 1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

应用	Ab评论	说明
IHC-Fr	★★★★☆	1/1000. (see Abreview submitted by Greg Gibson) We recommend using Goat Anti-Rabbit IgG H&L (Cy3®) preadsorbed (ab6939) secondary antibody
WB	★★★★☆	1/1000. Detects a band of approximately 92 kDa. When using colorimetric substrates such as BCIP/NBT use at a 1/5000 dilution (for chemiluminescent substrates). Detects a band of approximately 92-95 kDa (pro-form) and 82kDa (active form) (Human samples). Mouse MMP9 is larger, and on SDS PAGE gels runs about 102-105 kDa. Dilution optimised using Chromogenic detection.
IP		Use at an assay dependent concentration.
ELISA	★★★★☆	Use at an assay dependent concentration.
ICC/IF	★★★★☆	1/500.
ICC		Use at an assay dependent concentration.
IHC-FoFr	★★★★★	Use at an assay dependent concentration. PubMed: 19295156

靶标

功能	May play an essential role in local proteolysis of the extracellular matrix and in leukocyte migration. Could play a role in bone osteoclastic resorption. Cleaves KiSS1 at a Gly-Leu bond. Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N-terminal one quarter fragments. Degrades fibronectin but not laminin or Pz-peptide.
组织特异性	Produced by normal alveolar macrophages and granulocytes.
疾病相关	Intervertebral disc disease Metaphyseal anadysplasia 2
序列相似性	Belongs to the peptidase M10A family. Contains 3 fibronectin type-II domains. Contains 4 hemopexin repeats.
结构域	The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.
翻译后修饰	Processing of the precursor yields different active forms of 64, 67 and 82 kDa. Sequentially processing by MMP3 yields the 82 kDa matrix metalloproteinase-9. N- and O-glycosylated.
细胞定位	Secreted, extracellular space, extracellular matrix.

图片



Western blot - Anti-MMP9 antibody (ab38898)

All lanes : Anti-MMP9 antibody (ab38898) at 2 µg/ml

Lane 1 : Natural human MMP9 protein (Proenzyme, monomer) ([ab157344](#))

Lane 2 : Recombinant Mouse MMP9 protein ([ab39309](#))


Lysates/proteins at 0.1 µg per lane.

Secondary

Infrared labelled goat anti-rabbit (green) at 1/20000 dilution

Performed under reducing conditions.

This blot was produced using a 4-12% Bis-Tris gel under the MOPS buffer system. The gel was run at 200V for 60 minutes before being transferred onto a nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour before being incubated with anti-MMP9 antibody (ab38898; 2 microgram per mL) overnight at 4°C. Antibody binding was detected using infrared labelled goat anti-rabbit (green) antibody (diluted 1:20000) for 1 hour at room temperature before imaging.




Western blot - Anti-MMP9 antibody (ab38898)

Anti-MMP9 antibody (ab38898) + Human
MMP9

Observed band size : 88,92 kDa

Ab38898 detects a band at 92 Kd (pro-form) and a band at 88 Kd (active form). Mouse MMP9 is slightly larger than human MMP9, and the antibody detects a band at about 105 Kd. It is recommend to concentrate samples by Gelatin-agarose affinity chromatography prior to Western blot usage. A recommended starting concentration for Western blots is 1:1000 when using colorimetric substrates such as BCIP/NBT, and 1:5000 for chemiluminescent substrates. Higher concentration of antibody may be needed for non-human samples.



Western blot - Anti-MMP9 antibody (ab38898)

All lanes : Anti-MMP9 antibody (ab38898) at 5 µg/ml

Lane 1 : Recombinant Human MMP9, His tagged (ab82955) at 0.1 µg

Lane 2 : U937 whole cell lysate at 100 µg

Lane 3 : U937 whole cell lysate - treated with PMA and Brefeldin (24 hour treatment) at 100 µg


Lane 4 : Raw 264.7 (Mouse) whole cell lysate at 100 µg

Lane 5 : Raw 264.7 (Mouse) whole cell lysate - treated with LPS (6 hour treatment, 1ug/mL) at 100 µg

Performed under reducing conditions.

ab38898 detects recombinant Human MMP9 running at ~85 kDa, and endogenous full-length MMP9 in LPS-stimulated cells at ~100 kDa. This antibody also detects a band at 90 kDa in U937 PMA-treated cells.

ab38898 was incubated at 5 ug/mL and ab8245 (loading control to GAPDH) was diluted at 0.1 ug/mL and both were incubated for 48 hours at 4°C. Blots were developed with goat anti-rabbit IgG (H + L) and goat anti-mouse IgG (H + L) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



All lanes : Anti-MMP9 antibody (ab38898) at 1/1000 dilution

Lane 1 : Human lung tissue lysate at 100 µg with 10% Milk for 1 hour at room temperature

Lane 2 : Human lung tissue lysate at 100 µg with 10% Milk for 1 hour at room temperature

Lane 3 : MMP-9 KO mice tissue lysates with 10% Milk for 1 hour at room temperature

Lane 4 : MMP-9 KO mice tissue lysates with 10% Milk for 1 hour at room temperature

Secondary


HRP-conjugated donkey anti-rabbit polyclonal at 1/1000 dilution

Developed using the ECL technique

Performed under reducing conditions.

This image is courtesy of an anonymous Abreview

Specific observed bands 95-100 kDa




Immunocytochemistry/ Immunofluorescence - Anti-MMP9 antibody (ab38898)

Image courtesy of an anonymous Abreview

ab38898 staining MMP9 (red) in Mouse Neutrophils and Monocytes cells by ICC/IF (Immunocytochemistry/immunofluorescence).

Cells were fixed with paraformaldehyde and permeabilized with 2%BSA + 0.2% tritonX100 in PBS. Samples were incubated with primary antibody (1/200 in 2%BSA + 0.2% tritonX100 in PBS) for 25 minutes at 23°C. An Alexa Fluor® 568-conjugated Donkey anti-rabbit IgG polyclonal (1/1000) was used as the secondary antibody. DAPI is stained blue




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MMP9 antibody

(ab38898)

This image is courtesy of an anonymous Abreview

ab38898 staining MMP9 in Mouse Pancreatic carcinoma tissue sections by IHC-P (formaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 1% BSA for 1 hour at room temperature. Antigen retrieval was by heat mediation in citric acid (pH6). Samples were incubated with primary antibody (1/100) in 1% Aurion BSA for 12 hours. An HRP-conjugated Donkey polyclonal to rabbit IgG (1/100) was used as secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MMP9 antibody

(ab38898)

Image from Jung IH et al., PLoS One. 2011;6(12):e27941. Epub 2011 Dec 2. Fig 7.; doi:10.1371/journal.pone.0027941; December 2, 2011, PLoS ONE 6(12): e27941.

ab38898 staining MMP9 in 6 month-old transgenic zebrafish pancreas (Ihha overexpression) by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections).

Sections were incubated with primary antibody (1/500) and HRP-conjugated secondary antibody colored using DAB solution. Slides were counterstained with hematoxylin.



Immunohistochemistry (Frozen sections) - Anti-MMP9 antibody (ab38898)

ab38898 at a 1/1000 dilution staining mouse heart tissue by Immunohistochemistry (Frozen sections). The tissue was removed from a mouse, rinsed in PBS and slowly frozen in supercooled isopentane. 14um sections were made using a cryostat. The sections were acetone fixed and blocked in 2% BSA prior to incubation with the MMP9 antibody. [Goat Anti-Rabbit IgG H&L \(Cy3®\) preadsorbed \(ab6939\)](#) was used as the secondary antibody. In the image: red staining = MMP9, blue staining = nuclei, green = gelatinase activity.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors