

Product datasheet

Anti-Myelin Basic Protein antibody [12] ab7349

★★★★☆ 11 Abreviews 72 References 4 图像

概述

产品名称	Anti-Myelin Basic蛋白抗体[12]
描述	大鼠单克隆抗体[12] to Myelin Basic蛋白
宿主	Rat
特异性	The antibody reacts weakly with peptides ending in the Phe 91 where the 91-92 Phe-Phe bond is broken. Synthetic peptide 82-99 reacts very well, as does intact MBP. Further epitope analysis indicates binding to a region defined by amino acids 82-87 (DENPVV). Mapped by Geysen method to DENPW method.
经测试应用	适用于: IHC-FoFr, IHC-P, WB, ELISA, RIA, IHC-Fr, ICC/IF
种属反应性	与反应: Mouse, Rat, Sheep, Rabbit, Guinea pig, Cow, Dog, Human, Pig, Apterionotus leptorhynchus
免疫原	Full length protein (Cow).
表位	Amino acids 82-87 (DENPVV).

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.1% Sodium Azide Constituents: Tissue Culture Supernatant
纯度	Tissue culture supernatant
克隆	单克隆
克隆编号	12
骨髓瘤	NS0
同种型	IgG2a

应用


Our [Abpromise guarantee](#) covers the use of **ab7349** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
IHC-FoFr	★★★★★	Use at an assay dependent concentration. PubMed: 18678887
IHC-P	★★★★★	Use at an assay dependent concentration.
WB	★★★★☆	Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration.
RIA		Use at an assay dependent concentration.
IHC-Fr	★★★★☆	Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration. PubMed: 23584610


靶标

功能	The classic group of MBP isoforms (isoform 4-isoform 14) are with PLP the most abundant protein components of the myelin membrane in the CNS. They have a role in both its formation and stabilization. The smaller isoforms might have an important role in remyelination of denuded axons in multiple sclerosis. The non-classic group of MBP isoforms (isoform 1-isoform 3/Golli-MBPs) may preferentially have a role in the early developing brain long before myelination, maybe as components of transcriptional complexes, and may also be involved in signaling pathways in T-cells and neural cells. Differential splicing events combined with optional post-translational modifications give a wide spectrum of isomers, with each of them potentially having a specialized function. Induces T-cell proliferation.
组织特异性	MBP isoforms are found in both the central and the peripheral nervous system, whereas Golli-MBP isoforms are expressed in fetal thymus, spleen and spinal cord, as well as in cell lines derived from the immune system.
疾病相关	Note=The reduction in the surface charge of citrullinated and/or methylated MBP could result in a weakened attachment to the myelin membrane. This mechanism could be operative in demyelinating diseases such as chronic multiple sclerosis (MS), and fulminating MS (Marburg disease).
序列相似性	Belongs to the myelin basic protein family.
发展阶段	Expression begins abruptly in 14-16 week old fetuses. Even smaller isoforms seem to be produced during embryogenesis; some of these persisting in the adult. Isoform 4 expression is more evident at 16 weeks and its relative proportion declines thereafter.
翻译后修饰	Several charge isomers of MBP; C1 (the most cationic, least modified, and most abundant form), C2, C3, C4, C5, C6, C7, C8-A and C8-B (the least cationic form); are produced as a result of optional PTM, such as phosphorylation, deamidation of glutamine or asparagine, arginine citrullination and methylation. C8-A and C8-B contain each two mass isoforms termed C8-A(H), C8-A(L), C8-B(H) and C8-B(L), (H) standing for higher and (L) for lower molecular weight. C3, C4 and C5 are phosphorylated. The ratio of methylated arginine residues decreases during aging, making the protein more cationic. The N-terminal alanine is acetylated (isoform 3, isoform 4, isoform 5 and isoform 6). Arg-241 was found to be 6% monomethylated and 60% symmetrically dimethylated.
细胞定位	Myelin membrane. Cytoplasmic side of myelin.



Anti-Myelin Basic Protein antibody [12]
(ab7349) + Mouse brain tissue lysate

Observed band size: 19,26 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Myelin Basic Protein antibody [12] (ab7349)

This image is courtesy of an Abreview submitted by April Rempel


ab7349 staining Myelin Basic Protein in human trigeminal ganglia tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and blocked with 10% serum for 1 hour at 20°C; antigen retrieval was by heat mediation in citrate buffer, pH 6.0. Samples were incubated with primary antibody (1/1000 in 10% NDS) for 16 hours at 22°C. [ab150165](#), a goat anti-rat IgG H&L (Alexa Fluor[®] 488) preadsorbed polyclonal (1/1000) was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Myelin Basic Protein antibody [12] (ab7349)

Image courtesy of an Abreview submitted by Guillermo Estivill-Torres.

ab7349 staining Myelin Basic Protein in murine cerebral cortex tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Tissue was not fixed. Samples were blocked with 2.5% serum for 1 hour followed by incubation with the primary antibody at a 1/200 dilution for 18 hours. A Biotin-conjugated rabbit anti-rat polyclonal was used as the secondary antibody at a 1/500 dilution.



Immunohistochemistry (Frozen sections) - Anti-Myelin Basic Protein antibody [12] (ab7349)

This image is courtesy of an Abreview submitted by Ms Nancy Nuttle-McMenemy

ab7349 at 1/100 staining mouse spinal cord tissue sections by IHC-Fr. The tissue was formaldehyde fixed and blocked with 3% serum prior to incubation with the antibody for 18 hours. A biotinylated horse anti-rat antibody was used as the secondary.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors