

N-Arachidonylglycine (NAGly) ab120346

1 图像

概述


产品名称	N-Arachidonylglycine (NAGly)
描述	T-type Ca ²⁺ channel blocker. Endogenous GLYT2抑制剂.
生物学描述	Endogenous inhibitor of GLYT2 with little or no effect on GLYT1 and is analgesic in rat models of neuropathic and inflammatory pain. Anandamide-like compound, with little affinity at CB ₁ receptors, VR1 receptors and anandamide transporters. Potent inhibitor of T-type Ca ²⁺ channels with little activity on anandamide-sensitive TRPV1 and TASK1 currents and high-voltage-activated (HVA) Ca _v 1.2, Ca _v 2.2, Na _v 1.7, and Na _v 1.8 currents.
纯度	> 98%

性能

化学名称 N-(1-Oxo-5Z,8Z,11Z,14Z-eicosatetraenyl)glycine

分子量 361.52

化学结构



分子式 C₂₂H₃₅NO₃

CAS编号 179113-91-8

PubChem识别号 5283389

存放说明 Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12 months.

溶解度概述 Soluble in ethanol to 100 mM


处理 Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.

Need more advice on solubility, usage and handling? Please visit our [frequently asked questions \(FAQ\) page](#) for more details.

SMILES O=C(CCC/C=CC/C=CC/C=CC/C=CCCC)NCC(=O)O

来源 Synthetic

图片



ab32557 staining p38 (phospho T180 + Y182) in RAW 264.7 cells treated with N-Arachidonylglycine (ab120346), by ICC/IF. Increase in expression of p38 (phospho T180 + Y182) correlates with increased concentration of N-Arachidonylglycine, as described in literature.

The cells were incubated at 37°C for 1h in media containing different concentrations of ab120346 (N-Arachidonylglycine) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab32557** (1/100 dilution) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors

- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team