

Product datasheet

Anti-PGRPS antibody [188C424] ab13903

★★★★★ 1 Abreviews 1 References 2 图像

概述

产品名称	Anti-PGRPS抗体[188C424]
描述	小鼠单克隆抗体[188C424] to PGRPS
宿主	Mouse
经测试应用	适用于: ICC/IF, WB
种属反应性	与反应: Mouse, Human
免疫原	Synthetic peptide: YVLKGHRDVQRTLSPG , corresponding to amino acids 165-180 of Human PGRPS. Run BLAST with ExPASy Run BLAST with NCBI
阳性对照	Tested with mouse spleen tissue.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: PBS, 0.05% BSA
纯度	Protein G purified
克隆	单克隆
克隆编号	188C424
同种型	IgG1

应用


Our [Abpromise guarantee](#) covers the use of **ab13903** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ICC/IF	★★★★★	


应用	Ab评论	说明
WB		
应用说明	WB: Use at a concentration of 2 µg/ml. Predicted molecular weight: 22 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.	
靶标		
功能	Pattern receptor that binds to murein peptidoglycans (PGN) of Gram-positive bacteria. Has bactericidal activity towards Gram-positive bacteria. May kill Gram-positive bacteria by interfering with peptidoglycan biosynthesis. Binds also to Gram-negative bacteria, and has bacteriostatic activity towards Gram-negative bacteria. Plays a role in innate immunity.	
组织特异性	Highly expressed in bone marrow. Weak expression found in kidney, liver, small intestine, spleen, thymus, peripheral leukocyte, lung, fetal spleen and neutrophils.	
序列相似性	Belongs to the N-acetylmuramoyl-L-alanine amidase 2 family.	
翻译后修饰	N-glycosylated. N-glycosylation is required for bactericidal activity.	
细胞定位	Secreted. Cytoplasmic granule.	

图片



ab13903 at 1/100 staining human bone Marrow derived CD34+ cells by ICC/IF. The cells were paraformaldehyde fixed, permeabilized and blocked with BSA before incubation with the antibody for 24 hours at 4°C. A FITC conjugated goat anti-mouse IgG was used as the secondary. The cells were counterstained with DAPI.

The image shows staining mainly in the cytoplasmic region.



Anti-PGRPS antibody [188C424] (ab13903)
at 2 µg/ml + Jurkat cell lysate

Predicted band size: 22 kDa

Observed band size: 28 kDa

Western blot - Anti-PGRPS antibody [188C424]
(ab13903)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors