

## Product datasheet

# Anti-PIM1 antibody [EP2645Y] ab75776

**重组** RabMAb

11 References 9 图像

### 概述

<b>产品名称</b>	Anti-PIM1抗体[EP2645Y]
<b>描述</b>	兔单克隆抗体[EP2645Y] to PIM1
<b>特异性</b>	Ab75776 does not recognize PIM1L (isoform 1, 44 kDa).
<b>经测试应用</b>	<b>适用于:</b> ICC/IF, WB, IHC-P, Flow Cyt <b>不适用于:</b> IP
<b>种属反应性</b>	<b>与反应:</b> Mouse, Human <b>预测可用于:</b> Rat
<b>免疫原</b>	A synthetic peptide corresponding to residues in human PIM1
<b>阳性对照</b>	Human prostatic adenocarcinoma tissue; LnCap, Jurkat and K562 cell lysates.
<b>常规说明</b>	<b>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</b>  This product is a recombinant rabbit monoclonal antibody.

### 性能

<b>形式</b>	Liquid
<b>存放说明</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
<b>存储溶液</b>	PBS 49%,Sodium azide 0.01%,Glycerol 50%,BSA 0.05%
<b>纯度</b>	Tissue culture supernatant
<b>克隆</b>	单克隆
<b>克隆编号</b>	EP2645Y
<b>同种型</b>	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab75776** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab 评论	说明
ICC/IF		Use a concentration of 5 µg/ml.
WB		1/10000 - 1/20000. Detects a band of approximately 45 kDa (predicted molecular weight: 45 kDa). Also detects a band of approximately 36kDa (isoform 2). The 45 kDa band is isoform 1.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
Flow Cyt		1/20.

**应用说明** Is unsuitable for IP.

## 靶标

**功能** May affect the structure or silencing of chromatin by phosphorylating HP1 gamma/CBX3. Isoform 2 promotes the G1/S transition of the cell cycle via up-regulation of CDK2 activity and phosphorylation of CDKN1B, resulting in enhanced nuclear export and proteasome-dependent degradation of CDKN1B. Isoform 2 also represses CDKN1B transcription by phosphorylating and inactivating the transcription factor FOXO3. Plays a role in signal transduction in blood cells. Contributes to both cell proliferation and survival and thus provides a selective advantage in tumorigenesis.


**组织特异性** Expressed primarily in cells of the hematopoietic and germline lineages. Isoform 1 and isoform 2 are both expressed in prostate cancer cell lines.

**序列相似性** Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. PIM subfamily. Contains 1 protein kinase domain.

**翻译后修饰** Autophosphorylated on both serine/threonine and tyrosine residues.

**细胞定位** Cytoplasm. Nucleus and Cell membrane.

## Anti-PIM1 antibody [EP2645Y] 图像



Western blot - Anti-PIM1 [EP2645Y] antibody (ab75776)

Anti-PIM1 antibody [EP2645Y] (ab75776) at 1/20000 dilution + Jurkat at 10 µg

### Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution


**Predicted band size** : 45 kDa

Blocking buffer and concentration

5% NFDN/TBST


Diluting buffer and concentration

5% NFDN /TBST




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - PIM1 antibody [EP2645Y] (ab75776)

ab75776 at 1/100-1/250 dilution staining PIM1 in human prostatic adenocarcinoma by Immunohistochemistry, Paraffin-embedded tissue.




Immunocytochemistry/ Immunofluorescence - Anti-PIM1 antibody [EP2645Y] (ab75776)

ICC/IF image of ab75776 stained HepG2 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab75776, 5µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.




Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling PIM1 with purified ab75776 at 1/20 dilution (10ug/mL) (red). Cells were fixed with 4% paraformaldehyde and permeabilised using 90% methanol. A goat anti rabbit IgG (Alexa Fluor®488) (ab150077) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (ab172730) (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody

Flow Cytometry - Anti-PIM1 antibody [EP2645Y] (ab75776)




ab75776 showing positive staining in Thyroid gland carcinoma tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)-Anti-PIM1 antibody [EP2645Y](ab75776)




ab75776 showing positive staining in Stomach adenocarcinoma tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)-Anti-PIM1 antibody [EP2645Y](ab75776)




ab75776 showing positive staining in Colonic adenocarcinoma tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)-Anti-PIM1 antibody [EP2645Y](ab75776)



ab75776 showing positive staining in Breast carcinoma tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)-Anti-PIM1 antibody [EP2645Y](ab75776)



ab75776 showing positive staining in Normal tonsil tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)-Anti-PIM1 antibody [EP2645Y](ab75776)

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

#### **Our Promise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors