


Product datasheet

Anti-PRPF8 antibody ab87433

2 References 2 图像

概述

产品名称	Anti-PRPF8抗体
描述	兔多克隆抗体to PRPF8
宿主	Rabbit
经测试应用	适用于: WB, ICC/IF
种属反应性	与反应: Mouse, Human 预测可用于: Rat, Cow 
免疫原	Synthetic peptide conjugated to KLH derived from within residues 2300 to the C-terminus of Human PRPF8.参阅Abcam的专有抗源政策(Peptide available as ab87434 .)
阳性对照	This antibody gave a positive signal in the following lysates: HeLa Whole Cell lysate; HeLa Nuclear Lysate; NIH 3T3 Whole Cell lysate; A431 Whole Cell Lysate; MEF1 Whole Cell Lysate.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab87433** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 250 kDa (predicted molecular weight: 274 kDa).

应用	Ab评论	说明
ICC/IF		Use a concentration of 1 µg/ml.

靶标

功能	Central component of the spliceosome, which may play a role in aligning the pre-mRNA 5'- and 3'-exons for ligation. Interacts with U5 snRNA, and with pre-mRNA 5'-splice sites in B spliceosomes and 3'-splice sites in C spliceosomes.
组织特异性	Widely expressed.
疾病相关	Defects in PRPF8 are the cause of retinitis pigmentosa type 13 (RP13) [MIM:600059]. RP leads to degeneration of retinal photoreceptor cells. Patients typically have night vision blindness and loss of midperipheral visual field. As their condition progresses, they lose their far peripheral visual field and eventually central vision as well. RP13 inheritance is autosomal dominant.
序列相似性	Contains 1 MPN (JAB/Mov34) domain.
结构域	The MPN domain has structural similarity with viral ribonucleases and RNase H, but unlike RNases, it does not bind any metal ions.
翻译后修饰	Phosphorylated upon DNA damage, probably by ATM or ATR.
细胞定位	Nucleus speckle.

图片



Western blot - Anti-PRPF8 antibody (ab87433)

All lanes : Anti-PRPF8 antibody (ab87433) at 1 µg/ml

Lane 1 : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 2 : HeLa (Human epithelial carcinoma cell line) Nuclear Lysate

Lane 3 : NIH 3T3 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

Lane 4 : A431 (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 5 : MEF1 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 274 kDa

Observed band size: 250 kDa

Additional bands at: 100 kDa, 175 kDa. We are unsure as to the identity of these extra bands.

The 250 kDa band observed is comparable to the molecular weight seen with other commercially available antibodies to PRPF8.



Immunocytochemistry/ Immunofluorescence -
PRPF8 antibody (ab87433)

ICC/IF image of ab87433 stained MCF-7 cells. The cells were 100% Methanol fixed (5 min) and then incubated in 1%BSA / 10% normal Goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab87433, 1 µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 Goat anti-Rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM. This antibody also gave a positive result in 100% Methanol fixed (5 min) Hek293, and HepG2 cells at 1µg/ml.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors