


Product datasheet

Anti-PSMC3IP antibody ab82772

1 图像

概述

产品名称	Anti-PSMC3IP抗体
描述	兔多克隆抗体to PSMC3IP
经测试应用	适用于: WB, ELISA
种属反应性	与反应: Human 预测可用于: Mouse, Rat, Rabbit, Goat, Horse, Chicken, Cow, Cat, Dog, Pig, Zebrafish 
免疫原	Synthetic peptide derived from within residues: PEEKEQVYRE RQKYCKEWRK RKRMTATELSD AILEGYPKSK KQFFEEVGIE , corresponding to internal sequence amino acids 143-192 of Human PSMC3IP (NP_037422)
阳性对照	721_B cell lysate

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: None Constituents: 2% Sucrose, PBS
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab82772** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 24 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.
ELISA		Use at an assay dependent concentration.

靶标

功能	Plays an important role in meiotic recombination. Stimulates DMC1-mediated strand exchange required for pairing homologous chromosomes during meiosis. The complex PSMC3IP/MND1 binds DNA, stimulates the recombinase activity of DMC1 as well as DMC1 D-loop formation from double-strand DNA. This complex stabilizes presynaptic RAD51 and DMC1 filaments formed on single strand DNA to capture double-strand DNA. This complex stimulates both synaptic and presynaptic critical steps in RAD51 and DMC1-promoted homologous pairing. May inhibit HIV-1 viral protein TAT activity and modulate the activity of proteasomes through association with PSMC3. Acts as a tissue specific coactivator of hormone-dependent transcription mediated by nuclear receptors.
组织特异性	Highly expressed in testis and colon.
疾病相关	Ovarian dysgenesis 3 (ODG3) [MIM:614324]: A disorder characterized by lack of spontaneous pubertal development, primary amenorrhea, uterine hypoplasia, and hypergonadotropic hypogonadism as a result of streak gonads. Note=The disease is caused by mutations affecting the gene represented in this entry.
序列相似性	Belongs to the HOP2 family.
翻译后修饰	PTM: Phosphorylated by PKA, PKC and MAPK.
细胞定位	Nucleus.

Anti-PSMC3IP antibody 图像



Anti-PSMC3IP antibody (ab82772) at 1 μ g/ml
+ 721_B cell lysate at 10 μ g

Secondary

HRP conjugated anti-Rabbit IgG at 1/50000 dilution

Predicted band size : 24 kDa

Observed band size : 24 kDa

Additional bands at : 28 kDa, 55 kDa. We are unsure as to the identity of these extra bands.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors