

Product datasheet

Anti-R Phycoerythrin antibody (FITC) ab34723

概述

产品名称	Anti-R Phycoerythrin荧光抗体(FITC)
描述	山羊多克隆抗体to R Phycoerythrin荧光(FITC)
宿主	Goat
偶联物	FITC. Ex: 493nm, Em: 528nm
特异性	This antibody cross reacts with B Phycoerythrin.
经测试应用	适用于: Immunomicroscopy, Flow Cyt
免疫原	Highly purified R Phycoerythrin from the seaweed <i>Gracilaria</i> .

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	Preservative: 0.01% Sodium Azide Constituents: 10mg/ml BSA, 0.15M Sodium chloride, 0.02M Potassium phosphate. pH 7.2
纯化说明	This antibody was prepared from monospecific antiserum by immunoaffinity chromatography using a R Phycoerythrin coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab34723** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
Immunomicroscopy		
Flow Cyt		

应用说明 Flow Cyt: Use at an assay dependent dilution.

IM: Use at an assay dependent dilution.

The antibody is also thought to be suitable for other antibody based fluorescent assays.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

靶标

相关性

Phycoerythrin is one of a series of fluorescent pigments known as phycobiliproteins, which are produced by red and blue green algae. It occurs in more than one form, and has found application in immunology and diagnostic medicine. B and R Phycoerythrins provide superior labeling compared to fluorescein and rhodamine, and are used for labeling antibodies, usually monoclonals. These dyes may also be coupled to enzymes and other proteins, nucleic acids, polypeptide hormones, drugs, etc. Since phycoerythrins absorb light maximally between 450 and 650nm they fill the need for an intense fluorescent dye in the longer wavelengths of the visible spectrum, thereby avoiding interference from naturally fluorescing biological substances. R Phycoerythrin (240 kDa) is a labile molecule that may dissociate into components upon exposure to reducing or denaturing agents.

细胞定位

Plastid; chloroplast; chloroplast thylakoid lumen. Periphery of the rods of the phycobilisome.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors