



Product datasheet

Anti-RNF8 antibody ab15850

★★★★☆ 2 Abreviews 2 图像

概述

产品名称	Anti-RNF8抗体
描述	山羊多克隆抗体to RNF8
经测试应用	适用于: ICC/IF, WB
种属反应性	与反应: Human
免疫原	Synthetic peptide: GEPGFFVTGDRAG-C , corresponding to N terminal amino acids 2-14 of Human RNF8. (Peptide available as <a href="#">ab23278</a> .)
	 <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a>
阳性对照	Human lung lysate.

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 0.5% BSA, 0.5mg/ml Tris saline. pH 7.3
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab15850** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
ICC/IF	★★★★☆	1/100.

应用	Ab评论	说明
WB	★★★★☆	Use a concentration of 0.1 - 1 µg/ml. Detects a band of approximately 55 kDa (predicted molecular weight: 56 kDa). Can be blocked with <a href="#">Human RNF8 peptide (ab23278)</a> .

## 靶标

<b>功能</b>	E3 ubiquitin-protein ligase required for assembly of repair proteins to sites of DNA damage. Catalyzes the 'Lys-63'-linked ubiquitination of histone H2A and H2AX. Following DNA double-strand breaks (DSBs), it is recruited to the sites of damage by ATM-phosphorylated MDC1, mediates the ubiquitination of histones H2A and H2AX, thereby promoting the formation of TP53BP1 and BRCA1 ionizing radiation-induced foci (IRIF). Promotes the formation of 'Lys-63'-linked polyubiquitin chains and functions with the specific ubiquitin-conjugating UBE2N/UBC13. Substrates that are polyubiquitinated at 'Lys-63' are usually not targeted for degradation. Enforces the G2/M DNA damage checkpoint. Controls the recruitment of UIMC1-BRCC3 (RAP80-BRCC36) and PAXIP1/PTIP to DNA damage sites following DNA double-strand breaks (DSBs). Ubiquitination of histone H2A requires UBE2N but not MMS2 (UBE2V2). May also ubiquitinate histone H2B. Catalyzes the 'Lys-63'-linked ubiquitination of PCNA. May be required for proper exit from mitosis after spindle checkpoint activation and may regulate cytokinesis. May play a role in the regulation of RXRA-mediated transcriptional activity. Not involved in RXRA ubiquitination by UBE2E2.
<b>组织特异性</b>	Ubiquitous. In fetal tissues, highest expression in brain, thymus and liver. In adult tissues, highest levels in brain and testis, lowest levels in peripheral blood cells.
<b>通路</b>	Protein modification; protein ubiquitination.
<b>序列相似性</b>	Belongs to the RNF8 family. Contains 1 FHA domain. Contains 1 RING-type zinc finger.
<b>发展阶段</b>	Low levels at the G1-S boundary increase in intensity during S phase and until the end of the G2 phase. Abruptly decreases in late mitosis (at protein level). Barely detectable in anaphase.
<b>结构域</b>	The FHA domain specifically recognizes and binds ATM-phosphorylated MDC1 and Thr-4827 phosphorylated HERC2.
<b>翻译后修饰</b>	Autoubiquitinated through 'Lys-48' and 'Lys-63' of ubiquitin. 'Lys-63' polyubiquitination is mediated by UBE2N. 'Lys-29'-type polyubiquitination is also observed, but it doesn't require its own functional RING-type zinc finger.
<b>细胞定位</b>	Nucleus. Midbody. Following DNA double-strand breaks, recruited to the sites of damage. During prophase, concomitant with nuclear envelope breakdown, localizes throughout the cell, with a dotted pattern. In telophase, again in the nucleus and also with a discrete dotted pattern in the cytoplasm. In late telophase and during cytokinesis, localizes in the midbody of the tubulin bridge joining the daughter cells. Does not seem to be associated with condensed chromosomes at any time during the cell cycle.


## 图片



Western blot - RNF8 antibody (ab15850)

Anti-RNF8 antibody (ab15850) at 0.1 µg/ml +  
Lysates prepared from human lung tissues at  
35 µg

**Predicted band size : 56 kDa**



Immunocytochemistry/ Immunofluorescence - Anti-  
RNF8 antibody (ab15850)

Image courtesy of Dr Andrew Cobb by Abreview.

ab15850 staining RNF8 in human U2OS cells  
by Immunocytochemistry/  
Immunofluorescence.

Cells were fixed in formaldehyde,  
permeabilized using NP40, blocked with 5%  
BSA for 1 hour at 25°C and then incubated with  
ab15850 at a 1/100 dilution. The secondary  
used was an Alexa-Fluor 488 conjugated  
donkey anti-goat polyclonal used at a 1/500  
dilution.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors