

Product datasheet

Ramos whole cell lysate ab3955

概述

产品名称

Ramos全细胞裂解物

常规说明

Cell line: Ramos (Burkitt's lymphoma).

Growth media: RPMI 1640 medium with 2 mM L-glutamine adjusted to contain 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES, 1.0 mM sodium pyruvate, and 10% heat-inactivated FBS.

Ramos cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylene diamine tetra acetic acid, 1 mM phenyl methyl sulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecyl sulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecyl sulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.

经测试应用

适用于: WB

性能

存放说明

Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.

存储溶液

Constituent: 5% Beta mercaptoethanol

纯度

Whole Cell Lysate

裂解物说明

Ramos cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylene diamine tetra acetic acid, 1 mM phenyl methyl sulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecyl sulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecyl sulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.

背景

Derived from a caucasian Burkitt's lymphoma which does not possess the Epstein Barr Virus (EBV) genome. EBV infectability and permanent conversion into EBV positive sub-lines is possible by in vitro infection. The cells have B lymphocyte characteristics, with surface associated mu and kappa chains. Cells are used as model of B lymphocytes and for apoptosis studies.

应用

Our [Abpromise guarantee](#) covers the use of **ab3955** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
----	------	----

WB Use at an assay dependent concentration. 10 µg to 20 µg per lane is recommended for mini gel.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors