

Product datasheet

Recombinant Human eIF4EBP2 protein ab104667

1 References 1 图像

概述

产品名称	Recombinant人eIF4EBP2 protein
蛋白长度	Full length protein

描述

性质	Recombinant
来源	Escherichia coli
氨基酸序列	
Accession	Q13542
种属	Human
序列	MGSSHHHHHSSGLVPRGSHMSSSAGSGHQPSQSRAIPTRTV AISDAAQL PHDYCTTPGGTLFSTTPGGTRIIYDRKFLDRRNSPMAQTTPCHLPNI PGVTSPGTLIEDSKVEVNNLNNLNNHDRKHAVGDDAQFEMDI
分子量	15 kDa including tags
氨基酸	1 to 120
标签	His tag N-Terminus

技术指标

Our [Abpromise guarantee](#) covers the use of **ab104667** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	SDS-PAGE Mass Spectrometry
质谱法	MALDI-TOF
纯度	> 85 % SDS-PAGE. purified by using anion-exchange chromatography (DEAE sepharose resin) and gel-filtration chromatography (Sephacryl S-200) with 20mM Tris pH 7.5, 2mM EDTA.
形式	Liquid

制备和贮存

稳定性和存储

Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Preservative: None

Constituents: 10% Glycerol, 0.1M Sodium chloride, 20mM Tris HCl, 1mM DTT, pH 8.0

常规信息

功能

Repressor of translation initiation involved in synaptic plasticity, learning and memory formation (By similarity). Regulates EIF4E activity by preventing its assembly into the eIF4F complex: hypophosphorylated form of EIF4EBP2 competes with EIF4G1/EIF4G3 and strongly binds to EIF4E, leading to repress translation. In contrast, hyperphosphorylated form dissociates from EIF4E, allowing interaction between EIF4G1/EIF4G3 and EIF4E, leading to initiation of translation (PubMed:25533957). EIF4EBP2 is enriched in brain and acts as a regulator of synapse activity and neuronal stem cell renewal via its ability to repress translation initiation (By similarity). Mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase and mTORC1 pathways.

序列相似性

Belongs to the eIF4E-binding protein family.

结构域


The TOS motif mediates interaction with RPTOR, leading to promote phosphorylation by mTORC1 complex.

Intrinsically disordered protein that undergoes folding upon phosphorylation (PubMed:25533957). Hypophosphorylated form interacts strongly with EIF4E using (1) the YXXXXLPhi motif, that undergoes a disorder-to-helix transition upon binding and (2) the secondary EIF4E binding sites (residues 78-82) (PubMed:24207126, PubMed:25533957). Phosphorylation at Thr-37 and Thr-46 induces folding of region encompassing residues from Pro-18 to Arg-62 of into a four-stranded beta-domain that sequesters the helical YXXXXLPhi motif into a buried beta-strand, blocking accessibility to EIF4E. Protein phosphorylated at Thr-37 and Thr-46 is however unstable and subsequent phosphorylation at Ser-65, Thr-70 and Ser-83 is required to stabilize the fold, decreasing affinity for EIF4E by a factor of 4000 (PubMed:24207126, PubMed:25533957).

翻译后修饰

Phosphorylation at Thr-37, Thr-46, Ser-65, Thr-70 and Ser-83 is mediated by MTOR and corresponds to the hyperphosphorylated form: it abolishes binding to EIF4E by inducing folding of intrinsically disordered regions (PubMed:24207126, PubMed:25533957). First phosphorylated at Thr-37 and Thr-46 by MTOR, inducing folding of region encompassing residues from Pro-18 to Arg-62 of into a four-stranded beta-domain that sequesters the helical YXXXXLPhi motif into a partly buried beta-strand, blocking accessibility to EIF4E. Protein phosphorylated at Thr-37 and Thr-46 is however unstable and subsequent phosphorylation at Ser-65, Thr-70 and Ser-83 is required to stabilize the fold, decreasing affinity for EIF4E by a factor of 4000 (PubMed:24207126, PubMed:25533957). Phosphorylated in response to insulin, EGF and PDGF.

Deamidated at Asn-99 and Asn-102 to aspartate (Asp) in brain. Deamidation promotes interaction with RPTOR, subsequent phosphorylation by mTORC1 and increased translation, leading to impair kinetics of excitatory synaptic transmission. Deamidation takes place during postnatal development, when the PI3K-Akt-mTOR signaling is reduced, suggesting it acts as a compensatory mechanism to promote translation despite attenuated PI3K-Akt-mTOR signaling in neuron development. Deamidation converts Asn residues into a mixture of Asp and isoaspartate; interactions with PCMT1 is required to prevent isoaspartate accumulation and convert isoaspartate to Asp.



15% SDS-PAGE showing ab104667 at approximately 15.1kDa (3 μ g).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors