

## Product datasheet

# Anti-Retinoid X Receptor gamma antibody ab53162

★★★★☆ 2 Abreviews 1 References 1 图像

### 概述

产品名称	Anti-Retinoid X Receptor gamma抗体
描述	兔多克隆抗体to Retinoid X Receptor gamma
宿主	Rabbit
特异性	ab53162 detects endogenous levels of total Retinoic X Receptor gamma protein.
经测试应用	适用于: WB, ELISA, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide derived from human Retinoic X Receptor gamma.
阳性对照	HepG2 cell extracts.

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), 150mM Sodium chloride, pH 7.4
纯度	Immunogen affinity purified
纯化说明	ab53162 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
克隆	多克隆
同种型	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab53162** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB	★★★★☆	1/500 - 1/1000. Detects a band of approximately 51 kDa (predicted molecular weight: 51 kDa).

应用	Ab评论	说明
ELISA		1/10000.
IHC-P		Use a concentration of 2 - 4 µg/ml.

## 靶标

### 功能

Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RAR/RXR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. The high affinity ligand for RXRs is 9-cis retinoic acid.

### 序列相似性

Belongs to the nuclear hormone receptor family. NR2 subfamily.  
Contains 1 nuclear receptor DNA-binding domain.


### 结构域

Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain.

### 细胞定位

Nucleus.

## 图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Retinoid X Receptor gamma antibody (ab53162)

Ab53162 staining Retinoid X Receptor gamma in human nucleus accumbens.

Left panel: with primary antibody at a dilution of 4 µg/ml. Right panel: isotype control.

Sections were stained using an automated system (DAKO Autostainer Plus), at room temperature: sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffers EDTA pH9.0 in a Dako PT link. Slides were peroxidase blocked in 3% H<sub>2</sub>O<sub>2</sub> in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors