

Product datasheet

Retinoid X Receptor gamma overexpression 293T lysate (whole cell) ab94313

2 图像

概述

| | |
|-------|--|
| 产品名称 | Retinoid X Receptor gamma overexpression 293T裂解液(全细胞) |
| 常规说明 | ab94313 is a 293T cell transfected lysate in which Human Retinoid X Receptor gamma has been transiently over-expressed using a pCMV-Retinoid X Receptor gamma plasmid. The lysate is provided in 1X Sample Buffer. Note: For more detailed how the transfected lysate was prepared view preparation notes |
| 经测试应用 | 适用于: WB |

性能


| | |
|------|---|
| 形式 | Liquid |
| 存放说明 | Shipped on dry ice. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| 存储溶液 | pH: 6.80 Constituent: 100% 1x Sample Buffer |
| 纯度 | Whole Cell Lysate |
| 背景 | Function: Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RAR/RXR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. The high affinity ligand for RXRs is 9-cis retinoic acid. Similarity: Belongs to the nuclear hormone receptor family. NR2 subfamily. Contains 1 nuclear receptor DNA-binding domain. Domain: Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain. |

应用


Our [Abpromise guarantee](#) covers the use of **ab94313** in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| 应用 | Ab评论 | 说明 |
|----|------|-------------------------------------|
| WB | | Use at an assay dependent dilution. |

Retinoid X Receptor gamma overexpression 293T lysate (whole cell) 图像



ab94313 at 15µg/lane on an SDS-PAGE gel



All lanes : Anti-Retinoid X Receptor gamma antibody ([ab57671](#)) at 1/500 dilution

Lane 1 : Retinoid X Receptor gamma overexpression 293T lysate (whole cell) (ab94313)

Lane 2 : 293T non-transfected lysate

Lysates/proteins at 25 µg per lane.

Secondary

Goat Anti-mouse IgG (H and L) HRP conjugated at 1/2500 dilution

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors