



Product datasheet

Anti-Thrombopoietin antibody ab40664

1 References

概述

产品名称	Anti-Thrombopoietin抗体
描述	兔多克隆抗体to Thrombopoietin
特异性	Cross reactivity: Thrombopoietin Human (299-332) 100%; Thrombopoietin Human recombinant ~90% Does not cross react with Erythropoietin Human recombinant.
经测试应用	适用于: ICC/IF, IHC-P, WB
种属反应性	与反应: Human 预测可用于: Chimpanzee 
免疫原	Synthetic peptide: PPTGQYTLFP LPPTLPTPVV QLHPLLDPDS APTP conjugated to KLH, corresponding to amino acids 299-332 of Human Thrombopoietin Peptide available as ab93626 . <div style="text-align: right;">  Run BLAST with  Run BLAST with </div>

性能

形式	Liquid
存放说明	Store at +4°C short term (1-2 weeks) (add antibacterial agent). Aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: None Constituents: Whole serum
纯度	Whole antiserum
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab40664** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ICC/IF		1/500.

IHC-P	Use at an assay dependent concentration. PubMed: 21149592
WB	1/800. Detects a band of approximately 59 kDa (predicted molecular weight: 38 kDa). Can be blocked with Human Thrombopoietin peptide (ab93626) . With this antibody, we have found that blocking with 5% goat or donkey serum significantly reduces background as compared to BSA or milk.

靶标

功能	Lineage-specific cytokine affecting the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may be the major physiological regulator of circulating platelets.
疾病相关	Defects in THPO are a cause of essential thrombocythemia (ET) [MIM:187950]. ET is inherited as an autosomal dominant trait which is characterized by elevated platelet levels due to sustained proliferation of megakaryocytes, and frequently lead to thrombotic and haemorrhagic complications.
序列相似性	Belongs to the EPO/TPO family.
结构域	Two-domain structure with an erythropoietin-like N-terminal and a Ser/Pro/Thr-rich C-terminal.
细胞定位	Secreted.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors