

Product datasheet

Anti-Transferrin antibody (FITC) ab34670

2 References

概述

产品名称	Anti-Transferrin抗体(FITC)
描述	兔多克隆抗体to Transferrin (FITC)
偶联物	FITC. Ex: 493nm, Em: 528nm
经测试应用	适用于: Immunomicroscopy, Flow Cyt
种属反应性	与反应: Human
免疫原	Transferrin from Human serum.
常规说明	4.5moles FITC (MW 390 Daltons) per mole of Rabbit IgG. Absorption Wavelength: 495 nm Emission Wavelength: 528 nm

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	Preservative: 0.01% (w/v) Sodium Azide Constituents: 10mg/ml Polyethylene glycol (PEG-8000), 0.15M Sodium chloride, 0.02M Potassium phosphate. pH 7.2
纯度	IgG fraction
纯化说明	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process including delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against buffer.
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab34670** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
Immunomicroscopy		
Flow Cyt		

应用	Ab评论	说明
----	------	----

应用说明

Flow Cyt: Use at an assay dependent dilution.
 IM: Use at an assay dependent dilution.
 Suitable for other antibody based fluorescent assays.

Not yet tested in other applications.
 Optimal dilutions/concentrations should be determined by the end user.

靶标

功能

Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in stimulating cell proliferation.

组织特异性

Expressed by the liver and secreted in plasma.

疾病相关

Defects in TF are the cause of atransferrinemia (ATRAF) [MIM:209300]. Atransferrinemia is rare autosomal recessive disorder characterized by iron overload and hypochromic anemia.

序列相似性

Belongs to the transferrin family.
 Contains 2 transferrin-like domains.

细胞定位

Secreted.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors