

Product datasheet

Anti-Vaccinia Virus antibody ab26853

1 图像

概述

产品名称	Anti-Vaccinia Virus抗体
描述	兔多克隆抗体to Vaccinia Virus
宿主	Rabbit
经测试应用	适用于: ELISA
种属反应性	Reacts with vaccinia virus.
免疫原	Synthetic peptide containing amino acids on the predicted N terminal of fusion protein (A27L) in vaccinia virus.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	Preservative: None Constituents: Whole serum.
纯度	Whole antiserum
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab26853** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ELISA		1/5000.

靶标

相关性

Vaccinia virus is an Orthopoxvirus, containing double stranded DNA. Fusion protein plays an important role in the entry of enveloped virus into cells. As vaccinia virus has a wide host range, it is conceivable that certain cellular components that are ubiquitously expressed on the cell mediate virus infection. The study of the entry process, attachment, fusion and the proteins and receptors involved is complex. During vaccinia virus infection, the fusion process is attributed to the action of the 14KDa protein (A27L). The N terminus of this protein recognises heparan sulfate on the cell surface. It interacts with the negative charges of sulfates of glycosaminoglycans (GAGs). Therefore, antibodies that recognize this 14KDa protein are able to neutralize vaccinia virus infection and enable identification other viral and cellular proteins which participate in the vaccinia virus entry process.

图片

Dilution	Pre-immune bleed	Purified polyclonal
100	0.272	3.276
1000	0.111	2.676
10000	0.064	1.048
100000	0.054	0.207
1000000	0.055	0.083

ELISA - Anti-Vaccinia Virus antibody (ab26853)

ELISA method:

Antigen is coated on EIA strips at 1µg per well.

Add 200µl blocking buffer then wash wells with PBST.

Antiserum or peptide specific purified antibody is diluted in series as 10²~10⁶ folds.

Incubate antibody for 1 hour.

Wash unbound antibodies and add HRP-conjugated anti-rabbit IgG.

Wash plates and add substrate to develop color for 5 minutes.

Read absorbance at 650nm. Amount of colour is directly proportional to amount of antibodies.

Antibody titer is defined as >0.1 of absorbance of antiserum minus pre-immune serum.

Titer ~ 1/437000.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors