

## Product datasheet

# Recombinant Human SIRT6 protein ab109044

### 概述

---

产品名称	Recombinant人SIRT6 protein
蛋白长度	Full length protein

### 描述

---

性质	Recombinant
来源	Escherichia coli

### 氨基酸序列

Accession	<a href="#">Q8N6T7</a>
种属	Human
分子量	45 kDa including tags
氨基酸	1 to 355
标签	His tag N-Terminus

### 技术指标

---

Our [Abpromise guarantee](#) covers the use of **ab109044** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	SDS-PAGE
内毒素水平	< 0.100 Eu/μg
纯度	> 90 % SDS-PAGE. ab109044 is 0.2 μm filtered.
形式	Liquid

### 制备和贮存

---

稳定性和存储	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. Preservative: None Constituents: 55mM Tris HCl, 150mM Sodium chloride, pH 8.2
--------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<b>功能</b>	NAD-dependent protein deacetylase. Has deacetylase activity towards histone H3K9Ac and H3K56Ac. Modulates acetylation of histone H3 in telomeric chromatin during the S-phase of the cell cycle. Deacetylates histone H3K9Ac at NF-kappa-B target promoters and may down-regulate the expression of a subset of NF-kappa-B target genes. Acts as a corepressor of the transcription factor HIF 1A to control the expression of multiple glycolytic genes to regulate glucose homeostasis. Required for genomic stability. Regulates the production of TNF protein. Has a role in the regulation of life span (By similarity). Deacetylation of nucleosomes interferes with RELA binding to target DNA. May be required for the association of WRN with telomeres during S-phase and for normal telomere maintenance. Required for genomic stability. Required for normal IGF1 serum levels and normal glucose homeostasis. Modulates cellular senescence and apoptosis. On DNA damage, promotes DNA end resection via deacetylation of RBBP8. Has very weak deacetylase activity and can bind NAD(+) in the absence of acetylated substrate.
<b>序列相似性</b>	Belongs to the sirtuin family. Class IV subfamily. Contains 1 deacetylase sirtuin-type domain.
<b>细胞定位</b>	Nucleus, nucleoplasm. Predominantly nuclear. Associated with telomeric heterochromatin regions.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors