

Product datasheet

Anti-CD45 antibody [LT40] (PE/Cy5®) ab25156

概述

| | |
|-------|--|
| 产品名称 | Anti-CD45抗体[LT40] (PE/Cy5®) |
| 描述 | 小鼠单克隆抗体[LT40] to CD45 (PE/Cy5®) |
| 宿主 | Mouse |
| 偶联物 | PE/Cy5®. Ex: 496nm, Em: 670nm |
| 经测试应用 | 适用于: Flow Cyt |
| 种属反应性 | 与反应: Chicken |
| 免疫原 | The details of the immunogen for this antibody are not available. |
| 常规说明 | This product or portions thereof is manufactured under license from Carnegie Mellon University under U.S. Patent Number 5,268,486 and related patents. Cy and CyDye are trademarks of GE Healthcare Limited. |

性能

| | |
|------|--|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at +4°C. |
| 存储溶液 | Preservative: 0.09% Sodium azide Constituents: PBS, 0.2% Glycine, 16% Sucrose |
| 纯度 | Stabilising agent. Purified IgM |
| 克隆 | 单克隆 |
| 克隆编号 | LT40 |
| 同种型 | IgM |
| 轻链类型 | kappa |

应用

Our [Abpromise guarantee](#) covers the use of **ab25156** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| 应用 | Ab评论 | 说明 |
|----------|------|--------------------------------------|
| Flow Cyt | | Use 0.2µg for 10 ⁶ cells. |

靶标

| | |
|--------------|--|
| 功能 | Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. |
| 疾病相关 | Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (T(-)B(+)NK(+)) SCID [MIM:608971]. A form of severe combined immunodeficiency (SCID), a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients present in infancy recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. Genetic variations in PTPRC are involved in multiple sclerosis susceptibility (MS) [MIM:126200]. MS is a neurodegenerative disorder characterized by the gradual accumulation of focal plaques of demyelination particularly in the periventricular areas of the brain. Peripheral nerves are not affected. Onset usually in third or fourth decade with intermittent progression over an extended period. The cause is still uncertain. |
| 序列相似性 | Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily. Contains 2 fibronectin type-III domains. Contains 2 tyrosine-protein phosphatase domains. |
| 结构域 | The first PTPase domain interacts with SKAP1. |
| 翻译后修饰 | Heavily N- and O-glycosylated. |
| 细胞定位 | Membrane. Membrane raft. Colocalized with DPP4 in membrane rafts. |

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors