

Product datasheet

Anti-Bu1a + Bu1b antibody [AV20] (Biotin) ab25166

概述

产品名称	Anti-Bu1a + Bu1b抗体[AV20] (Biotin)
描述	小鼠单克隆抗体[AV20] to Bu1a + Bu1b (Biotin)
宿主	Mouse
偶联物	Biotin
特异性	This antibody recognizes a monomorphic determinant on the Bu1 B cell associated alloantigens of both RPL 6 (Bu1a) and 7 (Bu1b) lines of inbred chickens. The antibody also identifies B cells in partially inbred birds.
经测试应用	适用于: Flow Cyt, IP, IHC-Fr
种属反应性	与反应: Chicken 不与反应: Turkey
免疫原	Tissue/ cell preparation: Bursal cells from one day old H.B15 (Bu1 ^{ab}) strain chickens.
常规说明	

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.1% Sodium Azide Constituents: PBS
纯度	IgG fraction
克隆	单克隆
克隆编号	AV20
同种型	IgG1
轻链类型	kappa

应用

Our [Abpromise guarantee](#) covers the use of **ab25166** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
Flow Cyt		Use 1µg for 10 ⁶ cells. ab18434 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IP		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration. Acetone fixed.

靶标

相关性

The Bu1 [chB6] molecule is expressed on chicken B cells throughout most of their development, as well as on some non lymphoid cells. It has long been used as an allotypic marker in important studies of B cell development, though its function is unknown. It has no recognizable similarity to known mammalian molecules and thus represents a unique B cell marker. Its presence in chickens may be related to differences in the properties of B cell development between chickens and mammalian species. The sequences of the different alleles of this gene revealed a higher level of polymorphism than expected. A restriction fragment length polymorphism linked to the CHB6 gene has been used to determine its location on the linkage map of the chicken genome, which will allow the definitive evaluation of reported associations with disease resistance. Bu1 is found on 85-90% of bursal cells, 2-8% of thymocytes, 15-27% of spleen cells, and 2-18% of peripheral blood cells. It is also expressed on subsets of macrophages and monocytes, but not on granulocytes, erythrocytes or thrombocytes.

细胞定位

Type I membrane protein.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors