

Product datasheet

Anti-Monocyte + Macrophage antibody [KUL01] ab25403

1 References

概述

产品名称	Anti-Monocyte + Macrophage抗体[KUL01]
描述	小鼠单克隆抗体[KUL01] to Monocyte + Macrophage
特异性	This antibody recognises monocytes and macrophages as well as interdigitating cells and activated microglia cells. The antibody does not react with B (Bu1 positive) or T (CD3 positive) lymphocytes.
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Chicken
免疫原	The details of the immunogen for this antibody are not available.
常规说明	This antibody is useful in the study of the development, distribution, function and ontogeny of the mononuclear phagocyte system (MPS) of the chicken by exclusively recognizing the cells of the MPS.

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	pH: 8.20 Constituent: 100% Borate buffered saline
纯度	IgG fraction
Primary antibody说明	This antibody is useful in the study of the development, distribution, function and ontogeny of the mononuclear phagocyte system (MPS) of the chicken by exclusively recognizing the cells of the MPS.
克隆	单克隆
克隆编号	KUL01
同种型	IgG1
轻链类型	kappa

应用

Our [Abpromise guarantee](#) covers the use of **ab25403** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
Flow Cyt		Use 1µg for 10 ⁶ cells. ab170190 -Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

靶标

相关性

Monocyte and macrophage are white blood cells that roam the body tissues engulfing foreign organisms. A monocyte is a leukocyte, part of the human body's immune system that protects against blood-borne pathogens and moves quickly (aprox. 8-12 hours) to sites of infection in the tissues. Monocytes are usually identified in stained smears by their large bi-lobed nucleus. Macrophages are cells within the tissues that originate from specific white blood cells called monocytes. Monocytes and macrophages are phagocytes, acting in both nonspecific defense (or innate immunity) as well as specific defense (or cell-mediated immunity) of vertebrate animals. Their role is to phagocytize (engulf and then digest) cellular debris and pathogens either as stationary or mobile cells, and to stimulate lymphocytes and other immune cells to respond to the pathogen.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors