

Product datasheet

Anti-UBE3A antibody ab58266

2 图像

概述

产品名称	Anti-UBE3A抗体
描述	小鼠单克隆抗体to UBE3A
宿主	Mouse
经测试应用	适用于: WB
种属反应性	与反应: Human
免疫原	Recombinant fragment: ETFQQLITYK VISNEFN SRN LVNDDDAVA ASKCLKMVYY ANVVGGEVDT NHNEEDDEEP IPESSELT LQ ELLGEE RRNK KGPRVDPLET ELGVKTLDCR , corresponding to amino acids 51-151 of Human UBE3A Run BLAST with ExPASy Run BLAST with NCBI

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: None PBS, pH 7.2
纯度	Protein G purified
克隆	单克隆
同种型	IgG2a
轻链类型	kappa

应用


Our [Abpromise guarantee](#) covers the use of **ab58266** in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB		Use a concentration of 1 - 5 µg/ml. Predicted molecular weight: 101 kDa.

靶标


功能	E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and transfers it to its substrates. Several substrates have been identified including the RAD23A and RAD23B, MCM7 (which is involved in DNA replication), annexin A1, the PML tumor suppressor, and the cell cycle regulator CDKN1B. Catalyzes the high-risk human papilloma virus E6-mediated ubiquitination of p53/TP53, contributing to the neoplastic progression of cells infected by these viruses. Additionally, may function as a cellular quality control ubiquitin ligase by helping the degradation of the cytoplasmic misfolded proteins. Finally, UBE3A also promotes its own degradation in vivo. Plays an important role in the regulation of the circadian clock: involved in the ubiquitination of the core clock component ARNTL/BMAL1, leading to its proteasomal degradation (PubMed:24728990).
通路	Protein modification; protein ubiquitination.
疾病相关	Angelman syndrome
序列相似性	Contains 1 HECT (E6AP-type E3 ubiquitin-protein ligase) domain.
翻译后修饰	Phosphorylation at Tyr-659 by ABL1 impairs E3 ligase activity and protects p53/TP53 from degradation in (HPV)-infected cells.
细胞定位	Nucleus. Cytoplasm.

图片



IHC-P - UBE3A antibody (ab58266)

UBE3A antibody (ab58266) used in immunohistochemistry at 3ug/ml on formalin fixed and paraffin embedded human lung.



Western blot - UBE3A antibody (ab58266)

UBE3A antibody (ab58266) at 1ug/lane + Jurkat cell lysate at 25ug/lane.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors