


Product datasheet

Anti-NIFK antibody ab61203

3 图像

概述

产品名称	Anti-NIFK抗体
描述	兔多克隆抗体to NIFK
宿主	Rabbit
特异性	ab61203 detects endogenous levels of total NIFK protein.
经测试应用	适用于: WB, ELISA, ICC/IF, IHC-P
种属反应性	与反应: Human 预测可用于: Mouse 
免疫原	Synthetic non-phosphopeptide derived from human NIFK around the phosphorylation site of threonine 234.
阳性对照	Human brain tissue and HuvEc cells.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS (without Mg ²⁺ and Ca ²⁺), 150mM Sodium chloride, pH 7.4
纯度	Immunogen affinity purified
纯化说明	ab61203 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab61203** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 42 kDa (predicted molecular weight: 34 kDa).
ELISA		1/5000.
ICC/IF		1/100 - 1/500.
IHC-P		1/50 - 1/100.

靶标

序列相似性

Contains 1 RRM (RNA recognition motif) domain.


翻译后修饰

Sequentially phosphorylated on Thr-238, Thr-234 and Ser-230. Thr-234 is phosphorylated only when Thr-238 is phosphorylated. Likewise, phosphorylation at Ser-230 requires that Thr-234 and Thr-238 are phosphorylated. Phosphorylation enhances MKI67 binding.

细胞定位

Nucleus > nucleolus. Chromosome. Localizes to mitotic chromosomes in conjunction with MKI67.


图片



ab61203 at 1/50 dilution staining NIFK in human brain by Immunohistochemistry, Paraffin-embedded tissue, in the absence (left) or presence (right) of the immunising peptide.



ab61203 at 1/100 dilution staining NIFK in HuvEc cells by Immunofluorescence, in the absence (left) or presence (right) of the immunising peptide.



All lanes : Anti-NIFK antibody (ab61203) at 1 µg/ml

Lane 1 : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 2 : SW480 (Human colon adenocarcinoma cell line) Whole Cell Lysate (ab76999) with SW480 (Human colon adenocarcinoma cell line) Whole Cell Lysate

Lane 3 : HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) (ab65484) at 1/3000 dilution

Predicted band size: 34 kDa

Observed band size: 41 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors