

Product datasheet

Anti-MCM7 antibody [47DC141 or DCS-141] (Biotin) ab79802

1 References

概述

产品名称	Anti-MCM7抗体[47DC141 or DCS-141] (Biotin)
描述	小鼠单克隆抗体[47DC141 or DCS-141] to MCM7 (Biotin)
偶联物	Biotin
经测试应用	适用于: WB, IP, IHC-P, ICC/IF
种属反应性	与反应: Mouse, Rat, Dog, Human, Xenopus laevis
免疫原	Recombinant full length Human MCM7 protein
阳性对照	MAD109, or PC12 cells. Tonsil or breast cancer tissue.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.09% Sodium Azide Constituents: 0.2% BSA, 10mM PBS, pH 7.4
纯度	Protein G purified
克隆	单克隆
克隆编号	47DC141 or DCS-141
同种型	IgG1

应用

Our [Abpromise guarantee](#) covers the use of **ab79802** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB		
IP		

应用	Ab评论	说明
IHC-P		
ICC/IF		
应用说明	<p>ICC/IF: Use at an assay dependent dilution.</p> <p>IHC-P: Use at a concentration of 0.5 - 1.0 µg/ml for 30 min at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.</p> <p>IP: Use at 2µg/mg of lysate. Native and denatured.</p> <p>WB: Use at a concentration of 1 - 2 µg/ml for 2hrs at RT. Predicted molecular weight: 81 kDa.</p> <p>Not yet tested in other applications.</p> <p>Optimal dilutions/concentrations should be determined by the end user.</p>	
靶标		
功能	<p>Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for S-phase checkpoint activation upon UV-induced damage.</p>	
序列相似性	<p>Belongs to the MCM family.</p> <p>Contains 1 MCM domain.</p>	
翻译后修饰	<p>Phosphorylated upon DNA damage, probably by ATM or ATR.</p>	
细胞定位	<p>Nucleus.</p>	

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors