

Product datasheet

Anti-CDK1 antibody [POH-1] ab8040

★★★★★ 2 Abreviews 4 References 4 图像

概述

产品名称	Anti-CDK1抗体[POH-1]
描述	小鼠单克隆抗体[POH-1] to CDK1
特异性	Reacts with Cdc2
经测试应用	适用于: ICC/IF, IP, WB, Flow Cyt, IHC-P
种属反应性	与反应: Mouse, Cow, Human, Mink, Monkey 不与反应: Rat, Saccharomyces cerevisiae, Xenopus laevis, Drosophila melanogaster, Schizosaccharomyces pombe
免疫原	Recombinant full length protein (Human).
阳性对照	HeLa.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 15mM Sodium Azide Constituents: PBS, pH 7.4
纯度	>95% by SDS-PAGE
克隆	单克隆
克隆编号	POH-1
同种型	IgG2a

应用

Our [Abpromise guarantee](#) covers the use of **ab8040** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ICC/IF		Use a concentration of 1 µg/ml.
IP		Use at an assay dependent concentration.


应用	Ab评论	说明
WB	★★★★★	Use at an assay dependent concentration. Predicted molecular weight: 34 kDa.
Flow Cyt		Use 1µg for 10 <sup>6</sup> cells. <a href="#">ab170191</a> -Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

## 靶标

功能	Plays a key role in the control of the eukaryotic cell cycle. It is required in higher cells for entry into S-phase and mitosis. p34 is a component of the kinase complex that phosphorylates the repetitive C-terminus of RNA polymerase II.
组织特异性	Isoform 2 is found in breast cancer tissues.
序列相似性	Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily. Contains 1 protein kinase domain.
细胞定位	Nucleus.

形式	CDK1 can be located to the Nucleus, cytoplasm and Mitochondria. It's cytoplasmic during interphase and reversibly translocated from cytoplasm to the nucleus when phosphorylated before G2-M transition when associated with cyclin-B1. Accumulates in mitochondria in G2-arrested cells upon DNA-damage.
----	---

## Anti-CDK1 antibody [POH-1] 图像




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Cdc2 antibody [POH-1] (ab8040)

Ab8040 staining human normal skin. Staining is localised to the cytoplasm.

Left panel: with primary antibody at 2 ug/ml.

Right panel: isotype control.


Sections were stained using an automated system DAKO Autostainer Plus , at room temperature. Sections were rehydrated and antigen retrieved with the Dako 3-in-1 antigen retrieval buffers EDTA pH 9.0 in a DAKO PT Link. Slides were peroxidase blocked in 3% H<sub>2</sub>O<sub>2</sub> in methanol for 10 minutes. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 minutes and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.



Immunocytochemistry/ Immunofluorescence - Cdc2 antibody [POH-1] (ab8040)

ICC/IF image of ab8040 stained MCF7 cells.

The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab8040, 1 µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-mouse IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Western blot - Anti-CDK1 antibody [POH-1] (ab8040)

Anti-CDK1 antibody [POH-1] (ab8040) at 1/3000 dilution + whole tissue lysate prepared from murine liver at 100 µg

### Secondary


HRP conjugated goat anti-mouse polyclonal  
Developed using the ECL technique

**Predicted band size :** 34 kDa

**Observed band size :** 34 kDa

**Exposure time :** 20 seconds

*Image courtesy of an anonymous Abreview*



Flow Cytometry-Anti-CDK1 antibody [POH-1] (ab8040)

Overlay histogram showing HeLa cells stained with ab8040 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab8040, 1µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (ab96879) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIGG2A] (ab91361, 1µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line). Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <http://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors