

Product datasheet

Anti-BDNF antibody ab88901

★★★★☆ 1 Abreviews 2 图像

概述

产品名称	Anti-BDNF抗体
描述	小鼠多克隆抗体to BDNF
经测试应用	适用于: WB
种属反应性	与反应: Mouse, Human
免疫原	Full length human BDNF protein (AAH29795).
阳性对照	NIH/3T3 cell lysate and BDNF transfected 293T cell lysate. Brain tissue lysate
常规说明	For BDNF, multiple WB bands are possible and expected. The human protein has 5 isoforms (precursors: 28 – 37 kDa) and can be glycosylated (Uniprot: http://www.uniprot.org/uniprot/P23560). The mature form is expected at ~14 kDa (monomer) and the dimer at ~28 kDa.

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: None Constituents: PBS, pH 7.2
纯度	Protein A purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab88901** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB	★★★★☆	1/500 - 1/1000. Predicted molecular weight: 28 kDa.

靶标

功能

During development, promotes the survival and differentiation of selected neuronal populations of the peripheral and central nervous systems. Participates in axonal growth, pathfinding and in the modulation of dendritic growth and morphology. Major regulator of synaptic transmission and plasticity at adult synapses in many regions of the CNS. The versatility of BDNF is emphasized by its contribution to a range of adaptive neuronal responses including long-term potentiation (LTP), long-term depression (LTD), certain forms of short-term synaptic plasticity, as well as homeostatic regulation of intrinsic neuronal excitability.

组织特异性

Brain. Highly expressed in hippocampus, amygdala, cerebral cortex and cerebellum. Also expressed in heart, lung, skeletal muscle, testis, prostate and placenta.

疾病相关

Bulimia nervosa 2
Congenital central hypoventilation syndrome

序列相似性

Belongs to the NGF-beta family.


翻译后修饰

The propeptide is N-glycosylated and glycosulfated.
Converted into mature BDNF by plasmin (PLG).

细胞定位

Secreted.


图片



Anti-BDNF antibody (ab88901) at 1/500 dilution + cell lysate from NIH/3T3 at 50 µg

Predicted band size : 28 kDa

Observed band size : 28 kDa



All lanes : Anti-BDNF antibody (ab88901) at 1/500 dilution

Lane 1 : cell lysate from BDNF transfected 293T cells

Lane 2 : cell lysate from non transfected 293T cells

Lysates/proteins at 25 µg per lane.

Predicted band size : 28 kDa

Observed band size : 28 kDa

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors