


Alsterpaullone, cyclin-dependent kinase (CDK) and GSK-3beta inhibitor ab141070


2 图像

概述

产品名称	Alsterpaullone, cyclin-dependent kinase (CDK) and GSK-3beta抑制剂
描述	Potent cyclin-dependent kinase (CDK) and GSK-3β抑制剂
纯度	> 96%
CAS编号	237430-03-4
化学结构	


性能

化学名称	7,12-Dihydro-9-nitroindolo[3,2- <i>d</i>][1]benzazepin-6(5 <i>H</i>)-one
分子量	293.28
分子式	C ₁₆ H ₁₁ N ₃ O ₃
PubChem识别号	5005498
存放说明	Store at +4°C. Store under desiccating conditions. The product can be stored for up to 12 months.
处理	<p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked questions (FAQ) page for more details.</p>
SMILES	[O-][N+](=O)c4cc3c(NC1C3CC(=O)Nc2ccccc12)cc4
来源	Synthetic



Chemical Structure - Alsterpaullone, cyclin-dependent kinase (CDK) and GSK-3beta inhibitor (ab141070)

2D chemical structure image of ab141070, Alsterpaullone, cyclin-dependent kinase (CDK) and GSK-3beta inhibitor



Immunocytochemistry/ Immunofluorescence - Alsterpaullone, cyclin-dependent kinase (CDK) and GSK-3beta inhibitor (ab141070)

ab32505 staining AKT1 in SK-N-SH cells treated with alsterpaullone (ab141070), by ICC/IF. Decrease of AKT1 expression correlates with increased concentration of alsterpaullone, as described in literature.

The cells were incubated at 37°C for 6h in media containing different concentrations of ab141070 (alsterpaullone) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab32505** (1/200 dilution) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team